arXiv:1504.05891v3 [cs.IT] 26 Apr 2016

Exponent Function for One Helper Source Coding Problem at Rates outside the Rate Region

Yasutada Oohama

***Abstract*—We consider the one helper source coding problem posed and investigated by Ahlswede, Körner and Wyner. Two correlated sources are separately encoded and are sent to a destination where the decoder wishes to decode one of the two sources with an arbitrary small error probability of decoding. In this system, the error probability of decoding goes to one as the source block length n goes to infinity. This implies that we have a strong converse theorem for the one helper source coding problem. In this paper we provide the much stronger version of this strong converse theorem for the one helper source coding problem. We prove that the error probability of decoding tends to one exponentially and derive an explicit lower bound of this exponent function.**

***Index Terms*—One helper source coding problem, strong converse theorem, exponent of correct probability of decoding**

I. INTRODUCTION

We consider the one helper source coding problem posed and investigated by Ahlswede, Körner and Wyner. Two correlated sources are separately encoded and are sent to a destination where the decoder wishes to decode one of the two sources with an arbitrary small error probability of decoding. In this system, the error probability of decoding goes to one as the source block length n goes to infinity. This implies that we have a strong converse theorem for the one helper source coding problem. In this paper we provide the much stronger version of this strong converse theorem for the one helper source coding problem. We prove that the error probability of decoding tends to one exponentially and derive an explicit lower bound of this exponent function.

II. PROBLEM FORMULATION

Let $\mathcal{X}$ and $\mathcal{Y}$ be finite sets and $\{(X_t,Y_t)\}_{t=1}^{\infty}$ be a stationary discrete memoryless source. For each $t=1,2,\cdots$, the random pair (X_t,Y_t) takes values in $\mathcal{X}\times\mathcal{Y}$, and has a probability distribution

$$p_{XY}=\{p_{XY}(x,y)\}_{(x,y)\in\mathcal{X}\times\mathcal{Y}}$$

We write n independent copies of $\{X_t\}_{t=1}^{\infty}$ and $\{Y_t\}_{t=1}^{\infty}$, respectively as

$$X^n=X_1,X_2,\cdots,X_n \text{ and } Y^n=Y_1,Y_2,\cdots,Y_n.$$



Fig. 1. One helper source coding system.

We consider a communication system depicted in Fig. 1. Data sequences X^n and Y^n are separately encoded to $\varphi_1^{(n)}(X^n)$ and $\varphi_2^{(n)}(Y^n)$ and those are sent to the information processing center. At the center the decoder function $\psi^{(n)}$ observes $(\varphi_1^{(n)}(X^n),\varphi_2^{(n)}(Y^n))$ to output the estimation $\hat{Y}^n$ of Y^n. The encoder functions $\varphi_1^{(n)}$ and $\varphi_2^{(n)}$ are defined by

$$\left.\begin{array}{l}\varphi_1^{(n)}:\mathcal{X}^n\to\mathcal{M}_1=\{\,1,2,\cdots,M_1\,\}\,,\\ \varphi_2^{(n)}:\mathcal{Y}^n\to\mathcal{M}_2=\{\,1,2,\cdots,M_2\,\}\,,\end{array}\right\} \tag{1}$$

where for each $i=1,2$, $\|\varphi_i^{(n)}\|$ $(=M_i)$ stands for the range of cardinality of $\varphi_i^{(n)}$. The decoder function $\psi^{(n)}$ is defined by

$$\psi^{(n)}:\mathcal{M}_1\times\mathcal{M}_2\,\to\,\mathcal{Y}^n. \tag{2}$$

The error probability of decoding is

$$\mathrm{P}_{\mathrm{e}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})=\Pr\left\{\hat{Y}^n\neq Y^n\right\}, \tag{3}$$

where $\hat{Y}^n=\psi^{(n)}(\varphi_1^{(n)}(X^n),\varphi_2^{(n)}(Y^n))$. A rate pair (R_1,R_2) is ε*-achievable* if for any $\delta>0$, there exist a positive integer $n_0=n_0(\varepsilon,\delta)$ and a sequence of triples $\{(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})\}_{n\geq n_0}$ such that for $n\geq n_0$,

$$\begin{aligned}&\frac{1}{n}\log\|\varphi_i^{(n)}\|\leq R_i+\delta \text{ for } i=1,2,\\&\mathrm{P}_{\mathrm{e}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})\leq\varepsilon.\end{aligned}$$

For $\varepsilon\in(0,1)$, the rate region $\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})$ is defined by

$$\begin{aligned}&\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})\\&\triangleq\{\,(R_1,R_2):(R_1,R_2)\;\text{ is }\varepsilon\text{-achievable for }p_{XY}\,\}\,.\end{aligned}$$

Furthermore, define

$$\mathcal{R}_{\mathrm{AKW}}(p_{XY})\triangleq\bigcap_{\varepsilon\in(0,1)}\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY}).$$

We can show that the two rate regions $\mathcal{R}_{\mathrm{AKW}}(\varepsilon|\,p_{XY})$, $\varepsilon\in(0,1)$ and $\mathcal{R}_{\mathrm{AKW}}(p_{XY})$ satisfy the following property.

Property 1:

a) The regions $\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})$, $\varepsilon\in(0,1)$, and $\mathcal{R}_{\mathrm{AKW}}(p_{XY})$ are closed convex sets of $\mathbb{R}_+^2$, where

$$\mathbb{R}_+^2\triangleq\{(R_1,R_2):R_1\geq0,R_2\geq0\}.$$

Y. Oohama is with University of Electro-Communications, 1-5-1 Chofugaoka Chofu-shi, Tokyo 182-8585, Japan.

b) $\mathcal{R}_{\rm AKW}(\varepsilon|p_{XY})$ has another form using (n,ε)-rate region $\mathcal{R}_{\rm AKW}(n,\varepsilon|p_{XY})$, the definition of which is as follows. We set

$$\begin{aligned}&\mathcal{R}_{\rm AKW}(n,\varepsilon|p_{XY})=\{(R_1,R_2):\\&\ \text{There exists } (\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)}) \text{ such that}\\&\frac{1}{n}\log||\varphi_i^{(n)}||\leq R_i, i=1,2,\\&{\rm P}_{\rm e}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})\leq\varepsilon\}.\end{aligned}$$

Using $\mathcal{R}_{\rm AKW}(n,\ \varepsilon|p_{XY})$, $\mathcal{R}_{\rm AKW}(\varepsilon|p_{XY})$ can be expressed as

$$\mathcal{R}_{\rm AKW}(\varepsilon|p_{XY})={\rm cl}\left(\bigcup_{m\geq 1}\bigcap_{n\geq m}\mathcal{R}_{\rm AKW}(n,\varepsilon|p_{XY})\right).$$

Proof of this property is given in Appendix A. It is well known that $\mathcal{R}_{\rm AKW}(p_{XY})$ was determined by Ahlswede, Körner and Wyner. To describe their result we introduce an auxiliary random variable U taking values in a finite set $\mathcal{U}$. We assume that the joint distribution of (U,X,Y) is

$$p_{UXY}(u,x,y)=p_U(u)p_{X|U}(x|u)p_{Y|X}(y|x).$$

The above condition is equivalent to $U\leftrightarrow X\leftrightarrow Y$. Define the set of probability distribution $p=p_{UXY}$ by

$$\mathcal{P}(p_{XY})\stackrel{\triangle}{=}\{p_{UXY}:|\mathcal{U}|\leq|\mathcal{X}|+1, U\leftrightarrow X\leftrightarrow Y\}.$$

Set

$$\begin{aligned}\mathcal{R}(p)&\stackrel{\triangle}{=}\{(R_1,R_2):R_1,R_2\geq 0\,,\\&\qquad R_1\geq I_p(X;U), R_2\geq H_p(Y|U)\},\\\mathcal{R}(p_{XY})&\stackrel{\triangle}{=}\bigcup_{p\in\mathcal{P}(p_{XY})}\mathcal{R}(p).\end{aligned}$$

We can show that the region $\mathcal{R}(p_{XY})$ is a closed convex set of $\mathbb{R}_+^2$. The rate region $\mathcal{R}_{\rm AKW}(p_{XY})$ was determined by Ahlswede and Körner [1] and Wyner [2]. Their result is the following.

Theorem 1 (Ahlswede, Körner [1] and Wyner [2]):

$$\mathcal{R}_{\rm AKW}(p_{XY})=\mathcal{R}(p_{XY}).$$

On the converse coding theorem Ahlswede *et al.* [3] obtained the following.

Theorem 2 (Ahlswede et al. [3]):] For each fixed $\varepsilon\in(0,1)$, we have

$$\mathcal{R}_{\rm AKW}(\varepsilon|p_{XY})=\mathcal{R}(p_{XY}).$$

Gu and Effors [5] examined a speed of convergence for ${\rm P}_{\rm e}^{(n)}$ to tend to 1 as $n\to\infty$ by carefully checking the proof of Ahlswede *et al.* [3]. However they could not obtain a result on an explicit form of the exponent function with respect to the code length n.

Our aim is to find an explicit form of the exponent function for the error probability of decoding to tend to one as $n\to\infty$ when $(R_1,R_2)\notin\mathcal{R}_{\rm AKW}(p_{XY})$. To examine this quantity, we define the following quantity. Set

$$\begin{aligned}&{\rm P}_{\rm c}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})\stackrel{\triangle}{=}1-{\rm P}_{\rm e}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)}),\\&G^{(n)}(R_1,R_2|p_{XY})\\&\stackrel{\triangle}{=}\min_{\substack{(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)}):\\(1/n)\log\|\varphi_i^{(n)}\|\leq R_i,i=1,2}}\left(-\frac{1}{n}\right)\log{\rm P}_{\rm c}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)}).\\&G(R_1,R_2|p_{XY})\stackrel{\triangle}{=}\lim_{n\to\infty}G^{(n)}(R_1,R_2|p_{XY}),\\&\mathcal{G}(p_{XY})\stackrel{\triangle}{=}\{(R_1,R_2,G):G\geq G(R_1,R_2|p_{XY})\}.\end{aligned}$$

By time sharing we have that

$$\begin{aligned}&G^{(n+m)}\left(\frac{nR_1+mR_1'}{n+m},\frac{nR_2+mR_2'}{n+m}\middle|p_{XY}\right)\\&\leq\frac{nG^{(n)}(R_1,R_2|p_{XY})+mG^{(m)}(R_1',R_2'|p_{XY})}{n+m}.\end{aligned}\tag{4}$$

Choosing $R=R'$ in (4), we obtain the following subadditivity property on $\{G^{(n)}(R_1,R_2|p_{XY})\ \}_{n\geq 1}$:

$$\begin{aligned}&G^{(n+m)}(R_1,R_2|p_{XY})\\&\leq\frac{nG^{(n)}(R_1,R_2|p_{XY})+mG^{(m)}(R_1,R_2|p_{XY})}{n+m},\end{aligned}$$

from which we have that $G^{(n)}(R_1,R_2|p_{XY})$ exists and satisfies the following:

$$\lim_{n\to\infty}G^{(n)}(R_1,R_2|p_{XY})=\inf_{n\geq 1}G^{(n)}(R_1,R_2|p_{XY}).$$

The exponent function $G(R_1,R_2|p_{XY})$ is a convex function of (R_1,R_2). In fact, from (4), we have that for any $\alpha\in[0,1]$

$$\begin{aligned}&G(\alpha R_1+\bar{\alpha}R_1',\alpha R_2+\bar{\alpha}R_2'|p_{XY})\\&\leq\alpha G(R_1,R_2|p_{XY})+\bar{\alpha}G(R_1',R_2'|p_{XY}).\end{aligned}$$

The region $\mathcal{G}(p_{XY})$ is also a closed convex set. Our main aim is to find an explicit characterization of $\mathcal{G}(p_{XY})$. In this paper we derive an explicit outer bound of $\mathcal{G}\ (p_{XY})$ whose section by the plane $G=0$ coincides with $\mathcal{R}_{\rm AKW}(p_{XY})$.

III. MAIN RESULT

In this section we state our main result. We first explain that the region $\mathcal{R}(p_{XY})$ has two other expressions using the supporting hyperplane. We define two sets of probability distributions on $\mathcal{U}\ \times\mathcal{X}\ \times\mathcal{Y}$ by

$$\begin{aligned}\mathcal{P}_{\rm sh}(p_{XY})&\stackrel{\triangle}{=}\{p=p_{UXY}:|\mathcal{U}|\leq|\mathcal{X}|, U\leftrightarrow X\leftrightarrow Y\},\\\mathcal{Q}(p_{Y|X})&\stackrel{\triangle}{=}\{q=q_{UXY}:|\mathcal{U}|\leq|\mathcal{X}|, U\leftrightarrow X\leftrightarrow Y,\\&\qquad p_{Y|X}=q_{Y|X}\}.\end{aligned}$$

For $(\alpha,\mu)\in(0,1]\times[0,1]$, set

$$\begin{aligned}
&R^{(\mu)}(p_{XY})\\
&\stackrel{\triangle}{=} \max_{p\in\mathcal{P}_{\rm sh}(p_{XY})}\left\{\bar{\mu}I_p(X;U)+\mu H_p(Y|U)\right\},\\
&\tilde{R}^{(\alpha,\mu)}(p_{XY})\\
&\stackrel{\triangle}{=} \min_{q\in\mathcal{Q}(p_{Y|X})}\{(1+\mu)\bar{\alpha}D(q_X||p_X)\\
&\qquad\qquad +\alpha[\bar{\mu}I_q(X;U)+\mu H_q(Y|U)]\},\\
&\mathcal{R}_{\rm sh}(p_{XY})\\
&\stackrel{\triangle}{=} \bigcap_{\mu\in[0,1]}\{(R_1,R_2):\bar{\mu}R_1+\mu R_2\geq R^{(\mu)}(p_{XY})\},\\
&\tilde{\mathcal{R}}_{\rm sh}^{(\alpha)}(p_{XY})\\
&\stackrel{\triangle}{=} \bigcap_{\mu\in[0,1]}\left\{(R_1,R_2):\bar{\mu}R_1+\mu R_2\geq\right.\\
&\qquad\qquad\qquad\left.\frac{1}{\alpha}\tilde{R}^{(\alpha,\mu)}(p_{XY})\right\},\\
&\tilde{\mathcal{R}}_{\rm sh}(p_{XY})\stackrel{\triangle}{=}\bigcap_{\alpha\in(0,1]}\tilde{\mathcal{R}}_{\rm sh}^{(\alpha)}(p_{XY})\\
&\stackrel{\triangle}{=} \bigcap_{\substack{\alpha\in(0,1],\\ \mu\in[0,1]}}\left\{(R_1,R_2):\bar{\mu}R_1+\mu R_2\geq\right.\\
&\qquad\qquad\qquad\left.\frac{1}{\alpha}\tilde{R}^{(\alpha,\mu)}(p_{XY})\right\}.
\end{aligned}$$

For $\mathcal{R}\subseteq\mathbb{R}_+^2$, we set

$$\mathcal{R}-\kappa(1,1)\stackrel{\triangle}{=}\{(a-\kappa,b-\kappa)\in\mathbb{R}_+^2:(a,b)\in\mathcal{R}\}.$$

Then we have the following property.

Property 2:

a) The bound $|\mathcal{U}|\leq|\mathcal{X}|$ is sufficient to describe $R^{(\mu)}(\,p_{XY})$ and $\tilde{R}^{(\alpha,\mu)}(\,p_{XY})$.

b) For any p_{XY} we have

$$\mathcal{R}_{\rm sh}(p_{XY})=\mathcal{R}(p_{XY}). \tag{5}$$

Furthermore, for any $\alpha\in(0,\alpha_0]$, we have

$$\begin{aligned}
&R^{(\mu)}(p_{XY})-c_1\sqrt{\frac{\alpha}{\bar{\alpha}}}\log\left(c_2\frac{\bar{\alpha}}{\alpha}\right)\\
&\leq\frac{1}{\alpha}\tilde{R}^{(\alpha,\mu)}(p_{XY})\leq R^{(\mu)}(p_{XY}),
\end{aligned}\tag{6}$$

where

$$\left.\begin{aligned}
\alpha_0&=\alpha_0(|\mathcal{X}|,|\mathcal{Y}|)\stackrel{\triangle}{=}[8\log(|\mathcal{X}||\mathcal{Y}|)+1]^{-1},\\
c_1&=c_1(|\mathcal{X}|,|\mathcal{Y}|)\stackrel{\triangle}{=}\frac{3}{2}\sqrt{\log(|\mathcal{X}||\mathcal{Y}|)^2},\\
c_2&=c_2(|\mathcal{X}|,|\mathcal{Y}|)\stackrel{\triangle}{=}\frac{(|\mathcal{X}|^4|\mathcal{Y}|)^{\frac{2}{3}}}{\log(|\mathcal{X}||\mathcal{Y}|)^2}.
\end{aligned}\right\}\tag{7}$$

The two inequalities of (6) implies that for each $\alpha\in(0,\alpha_0]$,

$$\begin{aligned}
&\mathcal{R}_{\rm sh}(p_{XY})-c_1\sqrt{\frac{\alpha}{\bar{\alpha}}}\log\left(c_2\frac{\bar{\alpha}}{\alpha}\right)(1,1)\\
&\subseteq\tilde{\mathcal{R}}_{\rm sh}^{(\alpha)}(p_{XY})\subseteq\mathcal{R}_{\rm sh}(p_{XY}).
\end{aligned}$$

Hence we have

$$\tilde{\mathcal{R}}_{\rm sh}(p_{XY})=\mathcal{R}_{\rm sh}(p_{XY}). \tag{8}$$

Property 2 part a) is also stated as Lemma 9 in Appendix B. Proof of this lemma is given in this appendix. Proof of Property 2 part b) is given in Appendix C. For $(\alpha,\mu)\in[0,1]^2$, define

$$\begin{aligned}
&\omega_{q_X||p_X,q_{XY|U}}^{(\alpha,\mu)}(x,y|u)\\
&\stackrel{\triangle}{=}(1+\mu)\bar{\alpha}\log\frac{q_X(x)}{p_X(x)}\\
&\quad+\alpha\left[\bar{\mu}\log\frac{q_{X|U}(x|u)}{q_X(x)}+\mu\log\frac{1}{q_{Y|U}(y|u)}\right],\\
&f_{q_X||p_X,q_{XY|U}}^{(\alpha,\mu,\lambda)}(x,y|u)\\
&\stackrel{\triangle}{=}\exp\left\{-\lambda\omega_{q_X||p_X,q_{XY|U}}^{(\alpha,\mu)}(x,y|u)\right\},\\
&\Omega^{(\alpha,\mu,\lambda)}(q|p_{XY})\\
&\stackrel{\triangle}{=}-\log\mathrm{E}_q\left[\exp\left\{-\lambda\omega_{q_X||p_X,q_{XY|U}}^{(\alpha,\mu)}(X,Y|U)\right\}\right]\\
&=-\log\left[\sum_{u,x,y}q_U(u)q_{X|U}(x|u)q_{Y|X}(y|x)\right.\\
&\qquad\left.\times f_{q_X||p_X,q_{XY|U}}^{(\alpha,\mu,\lambda)}(x,y|u)\right],\\
&\Omega^{(\alpha,\mu,\lambda)}(p_{XY})\stackrel{\triangle}{=}\min_{q\in\mathcal{Q}(p_{Y|X})}\Omega^{(\alpha,\mu,\lambda)}(q|p_{XY}),\\
&F^{(\alpha,\mu,\lambda)}(\bar{\mu}R_1+\mu R_2|p_{XY})\\
&\stackrel{\triangle}{=}\frac{\Omega^{(\alpha,\mu,\lambda)}(p_{XY})-\alpha\lambda(\bar{\mu}R_1+\mu R_2)}{1+\lambda(1+\mu)},\\
&F(R_1,R_2|p_{XY})\\
&\stackrel{\triangle}{=}\sup_{(\alpha,\mu)\in[0,1]^2,\lambda>0}F^{(\alpha,\mu,\lambda)}(\bar{\mu}R_1+\mu R_2|p_{XY}).
\end{aligned}$$

We can show that the above functions and sets satisfy the following property.

Property 3:

a) The cardinality bound $|\mathcal{U}|\leq|\mathcal{X}|$ in $\mathcal{Q}(p_{Y|X})$ is sufficient to describe the quantity $\Omega^{(\alpha,\mu,\lambda)}(p_{XY})$.

b) Define a probability distribution $q^{(\lambda)}=q_{UXY}^{(\lambda)}$ by

$$\begin{aligned}
&q^{(\lambda)}(u,x,y)\\
&\stackrel{\triangle}{=}\frac{q(u,x,y)\exp\left\{-\lambda\omega_{q_X||p_X,q_{XY|U}}^{(\alpha,\mu)}(x,y|u)\right\}}{\mathrm{E}_q\left[\exp\left\{-\lambda\omega_{q_X||p_X,q_{XY|U}}^{(\alpha,\mu)}(X,Y|U)\right\}\right]}.
\end{aligned}$$

Then we have

$$\begin{aligned}
&\frac{\mathrm{d}}{\mathrm{d}\lambda}\Omega^{(\alpha,\mu,\lambda)}(q|p_{XY})\\
&\qquad=\mathrm{E}_{q^{(\lambda)}}\left[\omega_{q_X||p_X,q_{XY|U}}^{(\alpha,\mu)}(X,Y|U)\right],\\
&\frac{\mathrm{d}^2}{\mathrm{d}\lambda^2}\Omega^{(\alpha,\mu,\lambda)}(q|p_{XY})\\
&\qquad=-\mathrm{Var}_{q^{(\lambda)}}\left[\omega_{q_X||p_X,q_{XY|U}}^{(\alpha,\mu)}(X,Y|U)\right].
\end{aligned}$$

The second equality implies that $\Omega^{(\alpha,\mu,\lambda)}(q|p_{XY})$ is a concave function of $\lambda>0$.

c) Define

$$\begin{aligned}\rho &= \rho(p_{XY}) \\ &\overset{\triangle}{=} \max_{\substack{q \in \mathcal{Q}(p_{Y|X}), \\ (\alpha,\mu) \in [0,1]^2}} \mathrm{Var}_q \left[\omega^{(\alpha,\mu)}_{q_X || p_X, q_{XY|U}}(X,Y|U) \right].\end{aligned}$$

Since

$$0 \leq \left[\omega^{(\alpha,\mu)}_{q_X || p_X, q_{XY|U}}(x,y|u) \right]^2 < +\infty$$

for $(u,x,y) \in \mathcal{U} \times \mathcal{X} \times \mathcal{Y}$, we have $\rho(p_{XY}) < \infty$. Then for any $\lambda \in (0,1]$, we have

$$\begin{aligned}&\Omega^{(\alpha,\mu,\lambda)}(q|p_{XY}) \\ &\geq \lambda \mathrm{E}_q \left[\omega^{(\alpha,\mu)}_{q_X || p_X, q_{XY|U}}(X,Y|U) \right] - \frac{1}{2}\rho(p_{XY})\lambda^2.\end{aligned}$$

Specifically, we have

$$\Omega^{(\alpha,\mu,\lambda)}(p_{XY}) \geq \lambda \tilde{R}^{(\alpha,\mu)}(p_{XY}) - \frac{1}{2}\rho(p_{XY})\lambda^2.$$

d) For any $\delta > 0$, there exists a positive number $\nu = \nu(\delta, |\mathcal{X}|, |\mathcal{Y}|) \in (0,1]$ such that for every $\tau \in (0,\nu]$, the condition $(R_1 + \tau, R_2 + \tau) \notin \mathcal{R}(p_{XY})$ implies

$$F(R_1, R_2|p_{XY}) \geq \frac{\rho(p_{XY})}{2} \cdot g^2 \left(\tfrac{\tau^{3+\delta}}{2\rho(p_{XY})} \right) > 0,$$

where g is the inverse function of $\vartheta(a) \overset{\triangle}{=} a + a^2, a > 0$.

Property 3 part a) follows from Lemma 10 in Appendix B. Proof of this lemma is given in this appendix. Proofs of Property 3 parts b),c), and d) are given in Appendix D. Our main result is the following.

Theorem 3: For any $R_1, R_2 \geq 0$, any p_{XY}, and for any $(\varphi_1^{(n)}, \varphi_1^{(n)}, \psi^{(n)})$ satisfying

$$\frac{1}{n} \log ||\varphi_i^{(n)}|| \leq R_i, i = 1, 2,$$

we have

$$\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)}, \varphi_2^{(n)}, \psi^{(n)}) \leq 5 \exp\left\{ -nF(R_1, R_2|p_{XY}) \right\}. \quad (9)$$

It follows from Theorem 3 and Property 3 part d) that if (R_1, R_2) is outside the capacity region, then the error probability of decoding goes to one exponentially and its exponent is not below $F(R_1, R_2|p_{XY})$. It immediately follows from Theorem 3 that we have the following corollary.

Corollary 1:

$$\begin{aligned}&G(R_1, R_2|p_{XY}) \geq F(R_1, R_2|p_{XY}), \\ &\mathcal{G}(p_{XY}) \subseteq \overline{\mathcal{G}}(p_{XY}) \\ &= \left\{ (R_1, R_2, G) : G \geq F(R_1, R_2|p_{XY}) \right\}.\end{aligned}$$

Proof of Theorem 3 will be given in the next section. The exponent function at rates outside the rate region was derived by Oohama and Han [7] for the separate source coding problem for correlated sources [6]. The techniques used by them is a method of types [8], which is not useful to prove Theorem 3. Some novel techniques based on the information spectrum method introduced by Han [9] are necessary to prove this theorem.

From Theorem 3 and Property 3 part d), we have the following corollary, which provides an explicit outer bound of $\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})$ with an asymptotically vanishing deviation from $\mathcal{R}_{\mathrm{AKW}}(p_{XY}) = \mathcal{R}(p_{XY})$. The strong converse theorem established by Ahlswede *et al.* [3] immediately follows from this corollary. From Theorem 3 and Property 3 part d) we have the following corollary.

Corollary 2: For each fixed $\varepsilon \in (0,1)$ and for any $\delta > 0$, there exists a positive integer n_0 with

$$n_0 = n_0(\varepsilon, \delta, |\mathcal{X}|, |\mathcal{Y}|, \rho(p_{XY}))$$

such that for $n \geq n_0$, we have

$$\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY}) \subseteq \mathcal{R}(p_{XY}) - \kappa_n(1,1),$$

where

$$\kappa_n \overset{\triangle}{=} \left\{ \sqrt{\tfrac{8\rho(p_{XY})}{n} \log\left(\tfrac{5}{1-\varepsilon}\right)} + \tfrac{4}{n}\log\left(\tfrac{5}{1-\varepsilon}\right) \right\}^{\frac{1}{3+\delta}}.$$

It immediately follows from the above result that for any $\varepsilon \in (0,1)$, we have

$$\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY}) = \mathcal{R}_{\mathrm{AKW}}(p_{XY}) = \mathcal{R}(p_{XY}).$$

Proof of this corollary will be given in the next section.

IV. PROOF OF THE MAIN RESULT

Let (X^n, Y^n) be a pair of random variables from the information source. We set $S = \varphi_1^{(n)}(X^n)$. Joint distribution $p_{SX^nY^n}$ of (S, X^n, Y^n) is given by

$$p_{SX^nY^n}(s, x^n, y^n) = p_{S|X^n}(s|x^n) \prod_{t=1}^{n} p_{X_tY_t}(x_t, y_t).$$

It is obvious that $S \leftrightarrow X^n \leftrightarrow Y^n$. Then we have the following.

Lemma 1: For any $\eta > 0$ and for any $(\varphi_1^{(n)}, \varphi_2^{(n)}, \psi^{(n)})$ satisfying

$$\frac{1}{n} \log ||\varphi_i^{(n)}|| \leq R_i, i = 1, 2,$$

we have

$$\begin{aligned}&\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)}, \varphi_2^{(n)}, \psi^{(n)}) \leq p_{SX^nY^n} \Bigg\{ \\ &\quad 0 \geq \frac{1}{n} \log \frac{\hat{q}_{SX^nY^n}(S, X^n, Y^n)}{p_{SX^nY^n}(S, X^n, Y^n)} - \eta, && (10) \\ &\quad 0 \geq \frac{1}{n} \log \frac{q_{X^n}(X^n)}{p_{X^n}(X^n)} - \eta, && (11) \\ &\quad R_1 \geq \frac{1}{n} \log \frac{p_{X^n|S}(X^n|S)}{p_{X^n}(X^n)} - \eta, \\ &\quad R_2 \geq \frac{1}{n} \log \frac{1}{p_{Y^n|S}(Y^n|S)} - \eta \Bigg\} + 4\mathrm{e}^{-n\eta}. && (12)\end{aligned}$$

The probability distribution appearing in the right members of (12) have a property that we can select them arbitrary. In (10), we can choose any probability distribution $\hat{q}_{SX^nY^n}$ on $\mathcal{S} \times \mathcal{X}^n \times \mathcal{Y}^n$. In (11), we can choose any distribution q_{X^n} on $\mathcal{X}^n$.

Proof of this lemma is given in Appendix E. From Lemma 1 we obtain the following lemma.

Lemma 2: For any $\eta > 0$ and for any $(\varphi_1^{(n)},\ \varphi_2^{(n)}, \psi^{(n)})$ satisfying

$$\frac{1}{n}\log ||\varphi_i^{(n)}|| \leq R_i, i = 1, 2,$$

we have

$$\begin{aligned}
&\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)}, \varphi_2^{(n)}, \psi^{(n)}) \\
&\leq p_{SX^nY^n}\Biggl\{0 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{q_{X_t}(X_t)}{p_{X_t}(X_t)} - \eta, \\
&R_1 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{p_{X_t|SX^{t-1}}(X_t|S, X^{t-1})}{p_{X_t}(X_t)} - \eta, \\
&R_2 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{1}{p_{Y_t|SX^{t-1}Y^{t-1}}(Y_t|S, X^{t-1}, Y^{t-1})} - 2\eta\Biggr\} \\
&\quad +4\mathrm{e}^{-n\eta}.
\end{aligned}$$

Proof: In (10) in Lemma 1, we choose $\hat{q}_{SX^nY^n}$ having the form

$$\begin{aligned}
&\hat{q}_{SX^nY^n}(S, X^n, Y^n) \\
&= p_S(S)\prod_{t=1}^{n}\left\{p_{X_t|SX^{t-1}Y^t}(X_t|S, X^{t-1}, Y^t)\right. \\
&\qquad \left.\times\, p_{Y_t|SY^{t-1}}(Y_t|S, Y^{t-1})\right\}.
\end{aligned}$$

In (11) in Lemma 1, we choose q_{X^n} having the form

$$q_{X^n}(X^n) = \prod_{t=1}^{n} q_{X_t}(X_t).$$

We further note that

$$\begin{aligned}
\frac{p_{X^n|S}(X^n|S)}{p_{X^n}(X^n)} &= \prod_{t=1}^{n}\frac{p_{X_t|SX^{t-1}}(X_t|S, X^{t-1})}{p_{X_t}(X_t)}, \\
p_{Y^n|S}(Y^n|S) &= \prod_{t=1}^{n} p_{Y_t|SY^{t-1}}(Y_t|S, Y^{t-1}).
\end{aligned}$$

Then the bound (12) in Lemma 1 becomes

$$\begin{aligned}
&\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)}, \varphi_2^{(n)}, \psi^{(n)}) \leq p_{SX^nY^n}\Biggl\{ \\
&\quad 0 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{p_{Y_t|SY^{t-1}}(Y_t|S, Y^{t-1})}{p_{Y_t|SX^{t-1}Y^{t-1}}(Y_t|S, X^{t-1}, Y^{t-1})} - \eta, \\
&\quad 0 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{q_{X_t}(X_t)}{p_{X_t}(X_t)} - \eta, \\
&R_1 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{p_{X_t|SX^{t-1}}(X_t|S, X^{t-1})}{p_{X_t}(X_t)} - \eta, \\
&R_2 \geq \frac{1}{n}\sum_{t=1}^{n}\frac{1}{p_{Y_t|SY^{t-1}}(Y_t|S, Y^{t-1})} - \eta\Biggr\} + 4\mathrm{e}^{-n\eta} \\
&\leq p_{SX^nY^n}\Biggl\{0 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{q_{X_t}(X_t)}{p_{X_t}(X_t)} - \eta, \\
&R_1 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{p_{X_t|SX^{t-1}}(X_t|S, X^{t-1})}{p_{X_t}(X_t)} - \eta, \\
&R_2 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{1}{p_{Y_t|SX^{t-1}Y^{t-1}}(Y_t|S, X^{t-1}, Y^{t-1})} - 2\eta\Biggr\} \\
&\quad +4\mathrm{e}^{-n\eta},
\end{aligned}$$

completing the proof. ■

Lemma 3: Suppose that for each $t = 1, 2, \cdots, n$, the joint distribution $p_{SX^tY^t}$ of the random vector SX^tY^t is a marginal distribution of $p_{SX^nY^n}$. Then we have the following Markov chain:

$$SX^{t-1} \leftrightarrow X_t \leftrightarrow Y_t \tag{13}$$

or equivalently that $I(Y_t; SX^{t-1}|X_t) = 0$. Furthermore, we have the following Markov chain:

$$Y^{t-1} \leftrightarrow SX^{t-1} \leftrightarrow (X_t, Y_t) \tag{14}$$

or equivalently that $I(X_tY_t; Y^{t-1}|SX^{t-1}) = 0$. The above two Markov chains are equivalent to the following one long Markov chain:

$$Y^{t-1} \leftrightarrow SX^{t-1} \leftrightarrow X_t \leftrightarrow Y_t. \tag{15}$$

Proof of this lemma is given in Appendix F. For $t = 1, 2, \cdots, n$, let U_t be a random variable taking values in $\mathcal{U}_t = \mathcal{M}_1 \times \mathcal{X}^{t-1}$. Define U_t by $U_t \overset{\triangle}{=} (S, X^{t-1})$. From Lemmas 2 and 3, we have the following.

Lemma 4: For any $\eta > 0$ and for any $(\varphi_1^{(n)},\ \varphi_2^{(n)}, \psi^{(n)})$ satisfying

$$\frac{1}{n}\log ||\varphi_i^{(n)}|| \leq R_i, i = 1, 2,$$

we have

$$\begin{aligned}
&\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)}, \varphi_2^{(n)}, \psi^{(n)}) \\
&\leq p_{SX^nY^n}\Biggl\{0 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{q_{X_t}(X_t)}{p_{X_t}(X_t)} - \eta, \\
&R_1 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{p_{X_t|U_t}(X_t|U_t)}{p_{X_t}(X_t)} - \eta, \\
&R_2 \geq \frac{1}{n}\sum_{t=1}^{n}\log\frac{1}{p_{Y_t|U_t}(Y_t|U_t)} - 2\eta\Biggr\} + 4\mathrm{e}^{-n\eta}.
\end{aligned}$$

The following is well known as the Cramèr's bound in the large deviation principle.

Lemma 5: For any real valued random variable Z and any $\lambda > 0$, we have

$$\Pr\{Z \geq a\} \leq \exp\left[-\left(\lambda a - \log \mathrm{E}[\exp(\lambda Z)]\right)\right].$$

For for $(\alpha, \mu) \in [0, 1]^2$ and $\lambda > 0$, we define

$$\begin{aligned}
&\Omega^{(\alpha,\mu,\lambda)}(q_{X^n}, S|p_{XY}) \overset{\triangle}{=} -\log \mathrm{E}\Biggl[\prod_{t=1}^{n} 1 \\
&\quad \times \exp\left\{-\lambda\omega^{(\alpha,\mu)}_{q_{X_t}||p_{X_t}, p_{X_tY_t|U_t}}(X_tY_t|U_t)\right\}\Biggr] \\
&= -\log \mathrm{E}\left[\prod_{t=1}^{n}\frac{q_{X_t}^{(1+\mu)\bar{\alpha}\lambda}(X_t)}{p_{X_t}^{(1+\mu)\bar{\alpha}\lambda}(X_t)}\frac{p_{X_t}^{\alpha\bar{\mu}\lambda}(X_t)p_{Y_t|U_t}^{\alpha\mu\lambda}(Y_t|U_t)}{p_{X_t|U_t}^{\alpha\bar{\mu}\lambda}(X_t|U_t)}\right].
\end{aligned}$$

By Lemmas 4 and 5, we have the following proposition.

Proposition 1: For any $(\alpha, \mu) \in [0, 1]^2$, any $\lambda > 0$, and any $(\varphi_1^{(n)},\ \varphi_2^{(n)}, \psi^{(n)})$ satisfying

$$\frac{1}{n}\log ||\varphi_i^{(n)}|| \leq R_i, i = 1, 2,$$

we have

$$\begin{aligned}
&\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})\\
&\le 5\exp\biggl\{-n\left[1+\lambda(1+\mu)\right]^{-1}\\
&\quad\times\left[\frac{1}{n}\Omega^{(\alpha,\mu,\lambda)}(q_{X^n},S|p_{XY})-\alpha\lambda(\bar{\mu}R_1+\mu R_2)\right]\biggr\}.
\end{aligned}$$

Proof: By Lemma 4, for $(\alpha,\mu)\in[0,1]^2$ and $\lambda>0$, we have the following chain of inequalities:

$$\begin{aligned}
&\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})\\
&\le p_{SX^nY^n}\left\{0\ge(1+\mu)\bar{\alpha}\left[\frac{1}{n}\sum_{t=1}^n\log\frac{q_{X_t}(X_t)}{p_{X_t}(X_t)}-\eta\right],\right.\\
&\quad\alpha\bar{\mu}R_1\ge\frac{1}{n}\sum_{t=1}^n\log\frac{p^{\alpha}_{X_t|U_t}(X_t|U_t)}{p^{\alpha}_{X_t}(X_t)}-\alpha\bar{\mu}\eta,\\
&\quad\left.\alpha\mu R_2\ge\frac{1}{n}\sum_{t=1}^n\log\frac{1}{p^{\alpha\mu}_{Y_t|U_t}(Y_t|U_t)}-2\alpha\mu\eta\right\}+4\mathrm{e}^{-n\eta}\\
&\le p_{SX^nY^n}\biggl\{\alpha(\bar{\mu}R_1+\mu R_2)+(1+\mu)\eta\\
&\quad\ge-\frac{1}{n}\sum_{t=1}^n\log\left[\frac{q^{(1+\mu)\bar{\alpha}}_{X_t}(X_t)}{p^{(1+\mu)\bar{\alpha}}_{X_t}(X_t)}\frac{p^{\alpha\bar{\mu}}_{X_t}(X_t)p^{\alpha\mu}_{Y_t|U_t}(Y_t|U_t)}{p^{\alpha\bar{\mu}}_{X_t|U_t}(X_t|U_t)}\right]\biggr\}\\
&\quad+4\mathrm{e}^{-n\eta}\\
&=p_{SX^nY^n}\biggl\{\frac{1}{n}\sum_{t=1}^n\log\biggl[\frac{q^{(1+\mu)\bar{\alpha}}_{X_t}(X_t)}{p^{(1+\mu)\bar{\alpha}}_{X_t}(X_t)}\\
&\qquad\times\frac{p^{\alpha\bar{\mu}}_{X_t}(X_t)p^{\alpha\mu}_{Y_t|U_t}(Y_t|U_t)}{p^{\alpha\bar{\mu}}_{X_t|U_t}(X_t|U_t)}\biggr]\\
&\qquad\ge-\left[\alpha(\bar{\mu}R_1+\mu R_2)+(1+\mu)\eta\right]\biggr\}+4\mathrm{e}^{-n\eta}\\
&\overset{\text{(a)}}{\le}\exp\biggl[n\biggl\{\alpha\lambda(\bar{\mu}R_1+\mu R_2)+\lambda(1+\mu)\eta\\
&\qquad-\frac{1}{n}\Omega^{(\alpha,\mu,\lambda)}(q_{X^n},S|p_{XY})\biggr\}\biggr]+4\mathrm{e}^{-n\eta}. \qquad (16)
\end{aligned}$$

Step (a) follows from Lemma 5. We choose η so that

$$\begin{aligned}
-\eta&=\alpha\lambda(\bar{\mu}R_1+\mu R_2)+\lambda(1+\mu)\eta\\
&\quad-\frac{1}{n}\Omega^{(\alpha,\mu,\lambda)}(q_{X^n},S|p_{XY}). \qquad (17)
\end{aligned}$$

Solving (17) with respect to η, we have

$$\eta=\frac{\frac{1}{n}\Omega^{(\alpha,\mu,\lambda)}(q_{X^n},S|p_{XY})-\alpha\lambda(\bar{\mu}R_1+\mu R_2)}{1+\lambda(1+\mu)}.$$

For this choice of η and (16), we have

$$\begin{aligned}
&\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})\le 5\mathrm{e}^{-n\eta}\\
&=5\exp\biggl\{-n\left[1+\lambda(1+\mu)\right]^{-1}\\
&\quad\times\left[\frac{1}{n}\Omega^{(\alpha,\mu,\lambda)}(q_{X^n},S|p_{XY})-\alpha\lambda(\bar{\mu}R_1+\mu R_2)\right]\biggr\},
\end{aligned}$$

completing the proof. ■

Set

$$\begin{aligned}
&\underline{\Omega}^{(\alpha,\mu,\lambda)}(p_{XY})\\
&\triangleq\inf_{n\ge1}\min_{\substack{S\in\mathcal{M}_1:\\ S\leftrightarrow X^n\leftrightarrow Y^n}}\max_{\substack{q_{X^n}:q_{X^n}=\\ \prod_{t=1}^n q_{X_t}}}\frac{1}{n}\Omega^{(\alpha,\mu,\lambda)}(q_{X^n},S|p_{XY}).
\end{aligned}$$

By Proposition 1 we have the folloing corollary.

Corollary 3: For any $(\alpha,\mu)\in[0,1]^2$, any $\lambda>0$, and any $(\varphi_1^{(n)},\ \varphi_2^{(n)},\psi^{(n)})$ satisfying

$$\frac{1}{n}\log||\varphi_i^{(n)}||\le R_i, i=1,2,$$

we have

$$\begin{aligned}
&\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})\\
&\le 5\exp\left\{-n\left[\frac{\underline{\Omega}^{(\alpha,\mu,\lambda)}(p_{XY})-\alpha\lambda(\bar{\mu}R_1+\mu R_2)}{1+\lambda(1+\mu)}\right]\right\}.
\end{aligned}$$

We shall call $\underline{\Omega}^{(\alpha,\mu,\lambda)}(p_{XY})$ the communication potential. The above corollary implies that the analysis of $\underline{\Omega}^{(\alpha,\mu,\lambda)}(p_{XY})$ leads to an establishment of a strong converse theorem for the one helper source coding problem. In the following argument we drive an explicit lower bound of $\underline{\Omega}^{(\alpha,\mu,\lambda)}(p_{XY})$. For each $t=1,2,\cdots,n$, set $u_t=(s,x^{t-1})\in\mathcal{U}_t$. For each $t=1,2,\cdots,n$, define

$$\begin{aligned}
&f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t,y_t|u_t)\\
&\triangleq\frac{q^{(1+\mu)\bar{\alpha}\lambda}_{X_t}(x_t)}{p^{(1+\mu)\bar{\alpha}\lambda}_{X_t}(x_t)}\frac{p^{\alpha\bar{\mu}\lambda}_{X_t}(x_t)p^{\alpha\mu\lambda}_{Y_t|U_t}(y_t|u_t)}{p^{\alpha\bar{\mu}\lambda}_{X_t|U_t}(x_t|u_t)}.
\end{aligned}$$

By definition we have

$$\begin{aligned}
&\exp\left\{-\Omega^{(\alpha,\mu,\lambda)}(q_{X^n},S|p_{XY})\right\}\\
&=\sum_{s,x^n,y^n}p_{SX^nY^n}(s,x^n,y^n)\prod_{t=1}^n f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t,y_t|u_t).
\end{aligned}$$

For each $t=1,2,\cdots,n$, we define the probability distribution

$$p^{(\alpha,\mu,\lambda)}_{SX^tY^t}\triangleq\left\{p^{(\alpha,\mu,\lambda)}_{SX^tY^t}(s,x^t,y^t)\right\}_{(s,x^t,y^t)\in\mathcal{M}_1\times\mathcal{X}^t\times\mathcal{Y}^t}$$

by

$$\begin{aligned}
&p^{(\alpha,\mu,\lambda)}_{SX^tY^t}(s,x^t,y^t)\triangleq C_t^{-1}p_{SX^tY^t}(s,x^t,y^t)\\
&\times\prod_{i=1}^t f^{(\alpha,\mu,\lambda)}_{q_{X_i}||p_{X_i},p_{X_iY_i|U_i}}(x_i,y_i|u_i)
\end{aligned}$$

where

$$\begin{aligned}
C_t\triangleq&\sum_{s,x^t,y^t}p_{SX^tY^t}(s,x^t,y^t)\\
&\qquad\times\prod_{i=1}^t f^{(\alpha,\mu,\lambda)}_{q_{X_i}||p_{X_i},p_{X_iY_i|U_i}}(x_i,y_i)
\end{aligned}$$

are constants for normalization. For $t=1,2,\cdots,n$, define

$$\Phi_t^{(\alpha,\mu,\lambda)}\triangleq C_tC_{t-1}^{-1}, \qquad (18)$$

where we define $C_0=1$. Then we have the following lemma.

Lemma 6: For each $t = 1, 2, \cdots, n$, and for any $(s, x^t, y^t) \in \mathcal{M}_1 \times \mathcal{X}^t \times \mathcal{Y}^t$, we have

$$\begin{aligned}
&p^{(\alpha,\mu,\lambda)}_{SX^tY^t}(s, x^t, y^t) = (\Phi^{(\alpha,\mu,\lambda)}_t)^{-1} p^{(\alpha,\mu,\lambda)}_{SX^{t-1}Y^{t-1}}(s, x^{t-1}, y^{t-1}) \\
&\quad \times p_{X_tY_t|SX^{t-1}Y^{t-1}}(x_t, y_t|s, x^{t-1}, y^{t-1}) \\
&\quad \times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t, y_t|u_t).
\end{aligned}$$

Furthermore, we have

$$\begin{aligned}
&\Phi^{(\alpha,\mu,\lambda)}_t \\
&= \sum_{s,x^t,y^t} p^{(\alpha,\mu,\lambda)}_{SX^{t-1}Y^{t-1}}(s, x^{t-1}, y^{t-1}) \\
&\qquad \times p_{X_tY_t|SX^{t-1}Y^{t-1}}(x_t, y_t|s, x^{t-1}, y^{t-1}) \\
&\qquad \times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t, y_t|u_t). \qquad (19)
\end{aligned}$$

Proof of this lemma is given in Appendix G. Define

$$\begin{aligned}
&p^{(\alpha,\mu,\lambda)}_{U_t}(u_t) = p^{(\alpha,\mu,\lambda)}_{SX^{t-1}}(s, x^{t-1}) \\
&\triangleq \sum_{y^{t-1}} p^{(\alpha,\mu,\lambda)}_{SX^{t-1}Y^{t-1}}(s, x^{t-1}, y^{t-1}).
\end{aligned}$$

From Lemmas 3 and 6, we have the following lemma.

Lemma 7:

$$\begin{aligned}
&\Phi^{(\alpha,\mu,\lambda)}_t \\
&= \sum_{u_t,x_t,y_t} p^{(\alpha,\mu,\lambda)}_{U_t}(u_t) p_{X_t|U_t}(x_t|u_t) p_{Y_t|X_t}(y_t|x_t) \\
&\qquad \times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t, y_t|u_t). \qquad (20)
\end{aligned}$$

Proof: By Lemma 3, we have

$$\begin{aligned}
&p_{X_tY_t|SX^{t-1}Y^{t-1}}(x_t, y_t|s, x^{t-1}, y^{t-1}) \\
&= p_{X_t|SX^{t-1}Y^{t-1}}(x_t|s, x^{t-1}, y^{t-1}) \\
&\qquad \times p_{Y_t|SX^tY^{t-1}}(x_t|s, x^{t-1}, y^{t-1}) \\
&= p_{X_t|SX^{t-1}Y^{t-1}}(x_t|s, x^{t-1}) p_{Y_t|X_t}(y_t|x_t)
\end{aligned}$$

for $(s, x^t, y^t) \in \mathcal{S} \times \mathcal{X}^t \times \mathcal{Y}^t$ and for $t = 1, 2, \cdots, n$.

Then by Lemma 6, we have

$$\begin{aligned}
&\Phi^{(\alpha,\mu,\lambda)}_t \\
&= \sum_{s,x^t,y^t} p^{(\alpha,\mu,\lambda)}_{SX^{t-1}Y^{t-1}}(s, x^{t-1}, y^{t-1}) \\
&\qquad \times p_{X_tY_t|SX^{t-1}Y^{t-1}}(x_t, y_t|s, x^{t-1}, y^{t-1}) \\
&\qquad \times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t, y_t|u_t), \\
&= \sum_{s,x^t,y^t} p^{(\alpha,\mu,\lambda)}_{SX^{t-1}Y^{t-1}}(s, x^{t-1}, y^{t-1}) \\
&\qquad \times p_{X_t|SX^{t-1}}(x_t|s, x^{t-1}) p_{Y_t|X_t}(y_t|x_t) \\
&\qquad \times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t, y_t|u_t) \\
&= \sum_{s,x^t,y_t} p^{(\alpha,\mu,\lambda)}_{SX^{t-1}}(s, x^{t-1}) \\
&\qquad \times p_{X_t|SX^{t-1}}(x_t|s, x^{t-1}) p_{Y_t|X_t}(y_t|x_t) \\
&\qquad \times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t, y_t|u_t),
\end{aligned}$$

completing the proof. ■

We have the following.

Lemma 8:

$$\Omega^{(\alpha,\mu,\lambda)}(q_{X^n}, S|p_{XY}) = \sum_{t=1}^{n} \log \Phi^{(\alpha,\mu,\lambda)}_t. \qquad (21)$$

Proof: From (18) we have

$$\log \Phi^{(\alpha,\mu,\lambda)}_t = -\log C_t + \log C_{t-1}. \qquad (22)$$

Furthermore, by definition we have

$$\Omega^{(\alpha,\mu,\lambda)}(q_{X^n}, S|p_{XY}) = -\log C_n, C_0 = 1. \qquad (23)$$

From (22) and (23), (21) is obvious. ■

The following proposition is a mathematical core to prove our main result.

Proposition 2: For any $\alpha \geq 1, \mu, \lambda > 0$, we have

$$\underline{\Omega}^{(\alpha,\mu,\lambda)}(p_{XY}) \geq \Omega^{(\alpha,\mu,\lambda)}(p_{XY}).$$

Proof: Set

$$\begin{aligned}
\mathcal{Q}_n(p_{Y|X}) &\triangleq \{q = q_{UXY} : |\mathcal{U}| \leq |\mathcal{M}_1||\mathcal{X}^{n-1}||\mathcal{Y}^{n-1}|, \\
&\qquad q_{Y|X} = p_{Y|X}, U \leftrightarrow X \leftrightarrow Y\}, \\
\hat{\Omega}^{(\alpha,\mu,\lambda)}_n(p_{XY}) &\triangleq \min_{q \in \mathcal{Q}_n(p_{Y|X})} \Omega^{(\alpha,\mu,\lambda)}(q|p_{XY}).
\end{aligned}$$

Let U_t be random variables taking values in $\mathcal{M}_1 \times \mathcal{X}^{t-1}$. We choose q_{U_t} so that

$$q_{U_t}(u_t) = p^{(\alpha,\mu,\lambda)}_{U_t}(u_t) \qquad (24)$$

and choose $q_{X_tY_t|U_t}$ so that

$$q_{X_tY_t|U_t}(x_t, y_t|u_t) = p_{X_tY_t|U_t}(x_t, y_t|u_t). \qquad (25)$$

Furthermore we have

$$q_{Y_t|X_tU_t}(y_t|x_t, u_t) \overset{\text{(a)}}{=} p_{Y_t|X_t}(y_t|x_t) = q_{Y_t|X_t}(y_t|x_t). \qquad (26)$$

Step (a) follows from Lemma 3. The equation (26) imply that

$$q_t = q_{U_tX_tY_t} \in \mathcal{Q}_n(p_{Y|X}).$$

Hence we have the following chain of inequalities:

$$\begin{aligned}
&\Phi^{(\alpha,\mu,\lambda)}_t \\
&\overset{\text{(a)}}{=} \sum_{u_t,x_t,y_t} q_{U_t}(u_t) p_{X_t|U_t}(x_t|u_t) p_{Y_t|X_t}(y_t|x_t) \\
&\qquad \times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t, y_t|u_t) \\
&\overset{\text{(b)}}{=} \sum_{u_t,x_t,y_t} q_{U_t}(u_t) p_{X_t|U_t}(x_t|u_t) p_{Y_t|X_t}(y_t|x_t) \\
&\qquad \times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},q_{X_tY_t|U_t}}(x_t, y_t|u_t) \\
&= \exp\left\{\Omega^{(\alpha,\mu,\lambda)}(q_t|p_{XY})\right\} \\
&\overset{\text{(c)}}{\geq} \exp\left\{\hat{\Omega}^{(\alpha,\mu,\lambda)}_n(p_{XY})\right\} \overset{\text{(d)}}{=} \exp\left\{\Omega^{(\alpha,\mu,\lambda)}(p_{XY})\right\}. \qquad (27)
\end{aligned}$$

Step (a) follows from Lemma 7. Step (b) follows from (25). Step (c) follows from $q_t \in \mathcal{Q}_n(p_{Y|X})$ and the definition of $\hat{\Omega}^{(\alpha,\mu,\lambda)}_n(p_{XY})$. Step (d) follows from Lemma 10 in Appendix B. To prove this lemma we bound the cardinality $|\mathcal{U}|$ appearing in the definition of $\hat{\Omega}^{(\alpha,\mu,\lambda)}_n(p_{XY})$ to show that the bound $|\mathcal{U}| \leq |\mathcal{X}|$ is sufficient to describe $\hat{\Omega}^{(\alpha,\mu,\lambda)}_n(p_{XY})$. Proof of Lemma 10 is given in Appendix B. Hence we have the following:

$$
\begin{aligned}
&\max_{\substack{q_{X^n}:q_{X^n}=\\ \prod_{t=1}^{n} q_{X_t}}} \frac{1}{n}\Omega^{(\alpha,\mu,\lambda)}(q_{X^n},S|p_{XY})\\
&\geq \frac{1}{n}\Omega^{(\alpha,\mu,\lambda)}(q_{X^n},S|p_{XY}) \overset{\text{(a)}}{=} \frac{1}{n}\sum_{t=1}^{n}\log\Phi_t^{(\alpha,\mu,\lambda)}\\
&\overset{\text{(b)}}{\geq} \Omega^{(\alpha,\mu,\lambda)}(p_{XY}). \qquad (28)
\end{aligned}
$$

Step (a) follows from Lemma 8. Step (b) follows from (27). Since (28) holds fo any $n \geq 1$ and any $p_{SX^nY^n}$ satisfying $S \leftrightarrow X^n \leftrightarrow Y^n$, we have that for any $\alpha \geq 1$ and for any positive μ, λ,

$$
\underline{\Omega}^{(\alpha,\mu,\lambda)}(p_{XY}) \geq \Omega^{(\alpha,\mu,\lambda)}(p_{XY}).
$$

Thus, Proposition 2 is proved. ■

Proof of Theorem 3: We have the following:

$$
\begin{aligned}
&\frac{1}{n}\log\left\{\frac{5}{\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})}\right\}\\
&\overset{\text{(a)}}{\geq} \frac{\underline{\Omega}^{(\alpha,\mu,\lambda)}(p_{XY})-\alpha\lambda(\bar{\mu}R_1+\mu R_2)}{1+\lambda(1+\mu)}\\
&\overset{\text{(b)}}{\geq} \frac{\Omega^{(\alpha,\mu,\lambda)}(p_{XY})-\alpha\lambda(\bar{\mu}R_1+\mu R_2)}{1+\lambda(1+\mu)}\\
&= F^{(\alpha,\mu,\lambda)}(\bar{\mu}R_1+\mu R_2|p_{XY}).
\end{aligned}
$$

Step (a) follows from Corollary 3. Step (b) follows from Proposition 2. Since the above bound holds for any positive α, μ, and λ, we have

$$
\frac{1}{n}\log\left\{\frac{5}{\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})}\right\} \geq F(R_1,R_2|p_{XY}).
$$

Thus (9) in Theorem 3 is proved. ■

Proof of Corollary 2: To prove this corollary we use the following expression of $\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY)}$ stated in Property 4 part b):

$$
\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY}) = \mathrm{cl}\left(\bigcup_{m\geq 1}\bigcap_{n\geq m}\mathcal{R}_{\mathrm{AKW}}(n,\varepsilon|p_{XY})\right). \qquad (29)
$$

We assume that

$$
(R_1,R_2) \in \bigcup_{m\geq 1}\bigcap_{n\geq m}\mathcal{R}_{\mathrm{AKW}}(n,\varepsilon|p_{XY}). \qquad (30)
$$

Then there exists a positive integer $m_0 = m_0(\varepsilon)$ and some $\{(\varphi_1^{(n)}, \varphi_2^{(n)}, \psi^{(n)})\}_{n\geq m_0}$ such that for $n \geq m_0(\varepsilon)$, we have

$$
\begin{aligned}
&\frac{1}{n}\log||\varphi^{(n)}|| \leq R_i, i=1,2\\
&\mathrm{P}_{\mathrm{e}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)}) \leq \varepsilon.
\end{aligned}
$$

Then by Theorem 3, we have

$$
\begin{aligned}
1-\varepsilon &\leq \mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)})\\
&\leq 5\exp\left\{-nF(R_1,R_2|p_{XY})\right\} \qquad (31)
\end{aligned}
$$

for any $n \geq m_0(\varepsilon)$. Fix any $\delta > 0$. We take a positive number $\nu = \nu(\delta,|\mathcal{X}|,|\mathcal{Y}|) \in (0,1]$ appearing in Property 3 part d) and set

$$
\begin{aligned}
\kappa_n &= \left\{2\rho(p_{XY})\vartheta\left(\sqrt{\tfrac{2}{n\rho(p_{XY})}\log\left(\tfrac{5}{1-\varepsilon}\right)}\right)\right\}^{\frac{1}{3+\delta}} \qquad (32)\\
&= \left\{\sqrt{\tfrac{8\rho(p_{XY})}{n}\log\left(\tfrac{5}{1-\varepsilon}\right)} + \tfrac{4}{n}\log\left(\tfrac{5}{1-\varepsilon}\right)\right\}^{\frac{1}{3+\delta}}.
\end{aligned}
$$

Since g is an inverse function of ϑ, (32) is equivalent to

$$
g\left(\tfrac{\kappa_n^{3+\delta}}{2\rho(p_{XY})}\right) = \sqrt{\tfrac{2}{n\rho(p_{XY})}\log\left(\tfrac{5}{1-\varepsilon}\right)}. \qquad (33)
$$

We take a sufficiently large positive integer m_1 so that we have $\kappa_n < \nu$ for $n \geq m_1$. Set $n_0 = \max\{m_0, m_1\}$. We claim that for $n \geq n_0$, we have $(R_1+\kappa_n, R_2+\kappa_n) \in \mathcal{R}(p_{XY})$. To prove this claim we suppose that $(R_1+\kappa_{n^*}, R_2+\kappa_{n^*})$ does not belong to $\mathcal{R}(p_{XY})$ for some $n^* \geq n_0$. Since $\mathcal{R}(p_{XY})$ is a closed set, there exists a positive number $\tau > \kappa_{n^*}$ sufficiently close to κ_{n^*} such that

$$
\begin{aligned}
&\kappa_{n^*} < \tau \leq \nu,\\
&(R_1+\tau, R_2+\tau) \notin \mathcal{R}(p_{XY}).
\end{aligned}
$$

Then we have the following chain of inequalities:

$$
\begin{aligned}
&5\exp\left[-n^*F(R_1,R_2|p_{XY})\right]\\
&\overset{\text{(a)}}{\leq} 5\exp\left[-\frac{n^*\rho(p_{XY})}{2}\cdot g^2\left(\tfrac{\tau^{3+\delta}}{2\rho(p_{XY})}\right)\right]\\
&\overset{\text{(b)}}{<} 5\exp\left[-\frac{n^*\rho(p_{XY})}{2}\cdot g^2\left(\tfrac{\kappa_{n^*}^{3+\delta}}{2\rho(p_{XY})}\right)\right]\\
&\overset{\text{(c)}}{=} 5\exp\left[-\frac{n^*\rho(p_{XY})}{2}\frac{2}{n^*\rho(p_{XY})}\log\left(\frac{5}{1-\varepsilon}\right)\right]\\
&= 1-\varepsilon. \qquad (34)
\end{aligned}
$$

Step (a) follows from Property 3 part d). Step (b) follows from $\kappa_{n^*} < \tau$. Step (c) follows from from (33). The bound (34) contradicts (31). Hence we have $(R_1+\kappa_n, R_2+\kappa_n) \in \mathcal{R}(p_{XY})$ or equivalent to

$$
(R_1,R_2) \in \mathcal{R}(p_{XY}) - \kappa_n(1,1)
$$

for $n \geq n_0$. Recalling the first assumption (30) on (R_1,R_2), we obtain

$$
\bigcup_{m\geq 1}\bigcap_{n\geq m}\mathcal{R}_{\mathrm{AKW}}(n,\varepsilon|p_{XY}) \subseteq \mathcal{R}(p_{XY}) - \kappa_n(1,1). \qquad (35)
$$

Taking the closure of both sides of (35), using (29), and considering that $\mathcal{R}(p_{XY}) - \kappa_n(1,1)$ is a closed set, we have that for $n \geq n_0$,

$$
\begin{aligned}
&\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})\\
&= \mathrm{cl}\left(\bigcup_{m\geq 1}\bigcap_{n\geq m}\mathcal{R}_{\mathrm{AKW}}(n,\varepsilon|p_{XY})\right) \subseteq \mathcal{R}(p_{XY}) - \kappa_n(1,1),
\end{aligned}
$$

completing the proof. ■



Fig. 2. One helper source coding system investigated by Wyner.

V. ONE HELPER PROBLEM STUDIED BY WYNER

We consider a communication system depicted in Fig. 2. Data sequences X^n, Y^n, and Z^n, respectively are separately encoded to $\varphi_1^{(n)}(X^n)$, $\varphi_2^{(n)}(Y^n)$, and $\varphi_3^{(n)}(Z^n)$. The encoded data $\varphi_1^{(n)}(X^n)$ and $\varphi_2^{(n)}(Y^n)$ and are sent to the information processing center 1. The encoded data $\varphi_1^{(n)}(X^n)$ and $\varphi_3^{(n)}(Z^n)$ and are sent to the information processing center 2. At the center 1 the decoder function $\psi^{(n)}$ observes $(\varphi_1^{(n)}(X^n)$, $\varphi_2^{(n)}(Y^n))$ to output the estimation $\hat{Y}^n$ of Y^n. At the center 2 the decoder function $\phi^{(n)}$ observes $(\varphi_1^{(n)}(X^n)$, $\varphi_3^{(n)}(Z^n))$ to output the estimation $\hat{Z}^n$ of Z^n. The error probability of decoding is

$$\begin{aligned}&\mathrm{P}_{\mathrm{e}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\varphi_3^{(n)},\psi^{(n)},\phi^{(n)})\\&=\Pr\left\{\hat{Y}^n\neq Y^n \text{ or } \hat{Z}^n\neq Z^n\right\},\end{aligned}$$

where $\hat{Y}^n=\psi^{(n)}(\ \varphi_1^{(n)}(X^n),\varphi_2^{(n)}(Y^n))$ and $\hat{Z}^n=\psi^{(n)}(\ \varphi_1^{(n)}(X^n),\varphi_3^{(n)}(Z^n))$.

A rate triple (R_1,R_2,R_3) is ε-*achievable* if for any $\delta>0$, there exist a positve interger $n_0=n_0(\varepsilon,\delta)$ and a sequence of three encoders and two decoders functions $\{(\varphi_1^{(n)},\varphi_2^{(n)},\ \varphi_3^{(n)},\psi^{(n)},\ \phi^{(n)})\}_{n\geq n_0}$ such that for $n\geq n_0(\varepsilon,\delta)$,

$$\begin{aligned}&\frac{1}{n}||\varphi_i^{(n)}||\leq R_i+\delta \text{ for } i=1,2,3,\\&\mathrm{P}_{\mathrm{e}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\varphi_3^{(n)},\psi^{(n)},\phi^{(n)})\leq\varepsilon.\end{aligned}$$

The rate region $\mathcal{R}_{\mathrm{W}}(\varepsilon|p_{XYZ})$ is defined by

$$\begin{aligned}&\mathcal{R}_{\mathrm{W}}(\varepsilon|p_{XYZ})\stackrel{\triangle}{=}\{(R_1,R_2,R_3):\\&(R_1,R_2,R_3) \text{ is } \varepsilon\text{-achievable for } p_{XYZ}\}.\end{aligned}$$

Furthermore, define

$$\mathcal{R}_{\mathrm{W}}(p_{XYZ})\stackrel{\triangle}{=}\bigcap_{\varepsilon\in(0,1)}\mathcal{R}_{\mathrm{W}}(\varepsilon|p_{XYZ}).$$

We can show that the two rate regions $\mathcal{R}_{\mathrm{W}}(\varepsilon|\ p_{XYZ})$, $\varepsilon\in(0,1)$ and $\mathcal{R}_{\mathrm{W}}(p_{XYZ})$ satisfy the following property.

Property 4:

a) The regions $\mathcal{R}_{\mathrm{W}}(\varepsilon|p_{XYZ})$, $\varepsilon\in(0,1)$, and $\mathcal{R}_{\mathrm{W}}(\ p_{XYZ})$ are closed convex sets of $\mathbb{R}_+^3$.

b) We set

$$\begin{aligned}&\mathcal{R}_{\mathrm{W}}(n,\varepsilon|p_{XYZ})=\{(R_1,R_2,R_3):\\&\ \text{There exists } (\varphi_1^{(n)},\varphi_2^{(n)},\varphi_3^{(n)},\psi^{(n)}) \text{ such that}\\&\frac{1}{n}\log||\varphi_i^{(n)}||\leq R_i, i=1,2,3\\&\mathrm{P}_{\mathrm{e}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\varphi_3^{(n)},\psi^{(n)})\leq\varepsilon\},\end{aligned}$$

which is called the (n,ε)-rate region. Using $\mathcal{R}_{\mathrm{W}}(n,$ $\varepsilon|p_{XYZ})$, $\mathcal{R}_{\mathrm{W}}(\varepsilon|p_{XYZ})$ can be expressed as

$$\mathcal{R}_{\mathrm{W}}(\varepsilon|p_{XYZ})=\mathrm{cl}\left(\bigcup_{m\geq 1}\bigcap_{n\geq m}\mathcal{R}_{\mathrm{W}}(n,\varepsilon|p_{XYZ})\right).$$

It is well known that $\mathcal{R}_{\mathrm{W}}(p_{XYZ})$ was determined by Wyner. To describe their result we introduce an auxiliary random variable U taking values in a finite set $\mathcal{U}$. We assume that the joint distribution of (U,X,Y) is

$$p_{UXY}(u,x,y,z)=p_U(u)p_{X|U}(x|u)p_{YZ|X}(y,z|x).$$

The above condition is equivalent to $U\leftrightarrow X\leftrightarrow YZ$. Define the set of probability distribution on $\mathcal{U}\ \times\mathcal{X}\ \times\mathcal{Y}\ \times\mathcal{Z}$ by

$$\begin{aligned}\mathcal{P}(p_{XYZ})\stackrel{\triangle}{=}\{&p=p_{UXYZ}:|\mathcal{U}|\leq|\mathcal{X}|+2,\\&U\leftrightarrow X\leftrightarrow YZ\}.\end{aligned}$$

Set

$$\begin{aligned}\mathcal{R}(p)&\stackrel{\triangle}{=}\{(R_1,R_2,R_3):R_1,R_2,R_3\geq 0,\\&\quad R_1\geq I_p(X;U),R_2\geq H_p(Y|U),\\&\quad R_3\geq H_p(Z|U)\},\\\mathcal{R}(p_{XYZ})&\stackrel{\triangle}{=}\bigcup_{p\in\mathcal{P}(p_{XYZ})}\mathcal{R}(p).\end{aligned}$$

We can show that the region $\mathcal{R}(p_{XYZ})$ is a closed convex set of $\mathbb{R}_+^3$. The rate region $\mathcal{R}_{\mathrm{W}}(p_{XYZ})$ was determined by Wyner [2]. His result is the following.

Theorem 4 (Wyner [2]):

$$\mathcal{R}_{\mathrm{W}}(p_{XYZ})=\mathcal{R}(p_{XYZ}).$$

On the strong converse theorem Csiszár and Körner [8] obtained the following.

Theorem 5 (Csiszár and Körner [8]): For each fixed $\varepsilon\in(0,1)$, we have

$$\mathcal{R}_{\mathrm{W}}(\varepsilon|p_{XYZ})=\mathcal{R}(p_{XYZ}).$$

To examine a rate of convergence for the error probability of decoding to tend to one as $n\to\infty$ for $(R_1,R_2,R_3)\notin$

$\mathcal{R}_{\rm W}(p_{XYZ})$, we define the following quantity. Set

$$
\begin{aligned}
&\mathrm{P}_{\rm c}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\varphi_3^{(n)},\psi^{(n)},\phi^{(n)})\\
&\triangleq 1-\mathrm{P}_{\rm e}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\varphi_3^{(n)},\psi^{(n)},\phi^{(n)}),\\
&G^{(n)}(R_1,R_2,R_3|p_{XYZ})\\
&\triangleq \min_{\substack{(\varphi_1^{(n)},\varphi_2^{(n)},\varphi_3^{(n)},\\ \psi^{(n)},\phi^{(n)}):\\ (1/n)\log\|\varphi_i^{(n)}\|\\ \leq R_i, i=1,2,3}} \left(-\frac{1}{n}\right)\log \mathrm{P}_{\rm c}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\varphi_3^{(n)},\psi^{(n)},\phi^{(n)}),\\
&G(R_1,R_2,R_3|p_{XYZ}) \triangleq \lim_{n\to\infty} G^{(n)}(R_1,R_2,R_3|p_{XYZ}),\\
&\mathcal{G}(p_{XYZ})\\
&\triangleq \{(R_1,R_2,R_3,G): G\geq G(R_1,R_2,R_3|p_{XYZ})\}.
\end{aligned}
$$

By time sharing we have that

$$
\begin{aligned}
&G^{(n+m)}\left(\tfrac{nR_1+mR_1'}{n+m},\tfrac{nR_2+mR_2'}{n+m},\tfrac{nR_2+mR_2'}{n+m}\middle| p_{XYZ}\right)\\
&\leq \frac{nG^{(n)}(R_1,R_2,R_3|p_{XYZ})+mG^{(m)}(R_1',R_2',R_3'|p_{XYZ})}{n+m}.
\end{aligned}
\tag{36}
$$

Choosing $R=R'$ in (36), we obtain the following subadditivity property on $\{G^{(n)}(R_1,R_2,R_3|p_{XYZ})\ \}_{n\geq 1}$:

$$
\begin{aligned}
&G^{(n+m)}(R_1,R_2,R_3|p_{XYZ})\\
&\leq \frac{nG^{(n)}(R_1,R_2,R_3|p_{XYZ})+mG^{(m)}(R_1,R_2,R_3|p_{XYZ})}{n+m},
\end{aligned}
$$

from which we have that $G(R_1,R_2,R_3|W_1,W_2)$ exists and satisfies the following:

$$
\begin{aligned}
&G(R_1,R_2,R_3|p_{XYZ})\\
&=\inf_{n\geq 1} G^{(n)}(R_1,R_2,R_3|p_{XYZ}).
\end{aligned}
$$

The exponent function $G(R_1,R_2,R_3|p_{XYZ})$ is a convex function of (R_1,R_2,R_3). In fact, by time sharing we have that

$$
\begin{aligned}
&G^{(n+m)}\left(\tfrac{nR_1+mR_1'}{n+m},\tfrac{nR_2+mR_2'}{n+m},\tfrac{nR_2+mR_2'}{n+m}\middle| p_{XYZ}\right)\\
&\leq \frac{nG^{(n)}(R_1,R_2,R_3|p_{XYZ})+mG^{(m)}(R_1',R_2',R_3'|p_{XYZ})}{n+m},
\end{aligned}
$$

from which we have that for any $\alpha\in[0,1]$

$$
\begin{aligned}
&G(\alpha R_1+\bar{\alpha}R_1',\alpha R_2+\bar{\alpha}R_2',\alpha R_3+\bar{\alpha}R_3'|p_{XYZ})\\
&\leq \alpha G(R_1,R_2,R_3|p_{XYZ})+\bar{\alpha}G(R_1',R_2',R_3'|p_{XYZ}).
\end{aligned}
$$

The region $\mathcal{G}(p_{XYZ})$ is also a closed convex set. Our main aim is to find an explicit characterization of $\mathcal{G}(p_{XYZ})$. In this paper we derive an explicit outer bound of $\mathcal{G}$ (p_{XYZ}) whose section by the plane $G=0$ coincides with $\mathcal{R}_{\rm W}(p_{XYZ})$. We first explain that the region $\mathcal{R}(p_{XYZ})$ has two other expressions using the supporting hyperplane. We define two sets of probability distributions on $\mathcal{U}\times\mathcal{X}\times\mathcal{Y}\times\mathcal{Z}$ by

$$
\begin{aligned}
\mathcal{P}_{\rm sh}(p_{XYZ}) &\triangleq \{p=p_{UXYZ}: |\mathcal{U}|\leq|\mathcal{X}|,\\
&\qquad U\leftrightarrow X\leftrightarrow YZ\},\\
\mathcal{Q}(p_{YZ|X}) &\triangleq \{q=q_{UXYZ}: |\mathcal{U}|\leq|\mathcal{X}|,\\
&\qquad p_{YZ|X}=q_{YZ|X}, U\leftrightarrow X\leftrightarrow YZ\}.
\end{aligned}
$$

For $(\alpha,\nu,\gamma)\in(0,1)\times[0,1]^2$, set

$$
\begin{aligned}
&R^{(\mu,\gamma)}(p_{XYZ})\\
&\triangleq \max_{p\in\mathcal{P}_{\rm sh}(p_{XYZ})} \{\bar{\mu}I_p(X;U)\\
&\qquad +\mu(\bar{\gamma}H_p(Y|U)+\gamma H_p(Z|U))\},\\
&\tilde{R}^{(\alpha,\mu,\gamma)}(p_{XYZ})\\
&\triangleq \min_{q\in\mathcal{Q}(p_{YZ|X})} \{(1+\mu)\bar{\alpha}D(q_X||p_X)+\alpha[\bar{\mu}I_q(X;U)\\
&\qquad +\mu(\bar{\gamma}H_q(Y|U)+\gamma H_q(Z|U))]\},\\
&\mathcal{R}_{\rm sh}(p_{XYZ})\\
&=\bigcap_{(\mu,\gamma)\in[0,1]^2}\{(R_1,R_2,R_3): \bar{\mu}R_1+\mu(\bar{\gamma}R_2+\gamma R_3)\\
&\qquad\qquad \geq R^{(\mu,\gamma)}(p_{XYZ})\},\\
&\tilde{\mathcal{R}}_{\rm sh}^{(\alpha)}(p_{XYZ})\\
&=\bigcap_{(\mu,\gamma)\in[0,1]^2}\biggl\{(R_1,R_2,R_3):\ \bar{\mu}R_1+\mu(\bar{\gamma}R_2+\gamma R_3)\\
&\qquad\qquad \geq \frac{1}{\alpha}\tilde{R}^{(\alpha,\mu,\gamma)}(p_{XYZ})\biggr\},\\
&\tilde{\mathcal{R}}_{\rm sh}(p_{XY}) \triangleq \bigcap_{\alpha\in(0,1]}\tilde{\mathcal{R}}_{\rm sh}^{(\alpha)}(p_{XYZ})\\
&=\bigcap_{\substack{\alpha\in(0,1],\\ (\mu,\gamma)\in[0,1]^2}}\biggl\{(R_1,R_2,R_3):\ \bar{\mu}R_1+\mu(\bar{\gamma}R_2+\gamma R_3)\\
&\qquad\qquad \geq \frac{1}{\alpha}\tilde{R}^{(\alpha,\mu,\gamma)}(p_{XYZ})\biggr\}.
\end{aligned}
$$

Then we have the following property.

Property 5:

a) The bound $|\mathcal{U}|\leq|\mathcal{X}|$ is sufficient to describe $R^{(\mu)}($ $p_{XYZ})$ and $\tilde{R}^{(\mu)}($ $p_{XYZ})$.

b) For any p_{XYZ} we have

$$
\mathcal{R}_{\rm sh}(p_{XYZ})=\mathcal{R}(p_{XYZ}). \tag{37}
$$

Furthermore, for any $\alpha\in(0,\alpha_0]$, we have

$$
\begin{aligned}
&R_{\rm sh}^{(\mu)}(p_{XYZ})-c_1\sqrt{\frac{\alpha}{\bar{\alpha}}}\log\left(c_2\frac{\bar{\alpha}}{\alpha}\right)\\
&\leq \frac{1}{\alpha}\tilde{R}_{\rm sh}^{(\alpha,\mu)}(p_{XYZ})\leq R_{\rm sh}^{(\mu)}(p_{XYZ}),
\end{aligned}
\tag{38}
$$

where $\alpha_0=\alpha_0(|\mathcal{X}|,|\mathcal{Y}|,|\mathcal{Z}|)$, $c_1=c_1(|\mathcal{X}|,|\mathcal{Y}|,|\mathcal{Z}|)$, and $c_2=c_2(|\mathcal{X}|,|\mathcal{Y}|,|\mathcal{Z}|)$ are some suitable constants. The two inequalities of (38) implies that for each $\alpha\in(0,\alpha_0]$,

$$
\begin{aligned}
&\mathcal{R}_{\rm sh}(p_{XYZ})-c_1\sqrt{\frac{\alpha}{\bar{\alpha}}}\log\left(c_2\frac{\bar{\alpha}}{\alpha}\right)(1,1,1)\\
&\subseteq \tilde{\mathcal{R}}_{\rm sh}^{(\alpha)}(p_{XYZ})\subseteq\mathcal{R}_{\rm sh}(p_{XYZ}).
\end{aligned}
$$

Hence we have

$$
\tilde{\mathcal{R}}_{\rm sh}(p_{XYZ})=\mathcal{R}_{\rm sh}(p_{XYZ}). \tag{39}
$$

Property 6:

$$
\mathcal{R}_{\rm sh}(p_{XYZ})=\tilde{\mathcal{R}}_{\rm sh}(p_{XYZ})=\mathcal{R}(p_{XYZ}).
$$

Since the proof of Property 6 is quite similar to that of Property 2, we omit it. For $(\alpha, \mu, \gamma) \in [0,1]^3$, define

$$\begin{aligned}
&\omega^{(\alpha,\mu,\gamma)}_{q_X||p_X,q_{XYZ|U}}(x,y,z|u) \\
&\triangleq (1+\mu)\bar{\alpha} \log \frac{q_X(x)}{p_X(x)} + \alpha \Biggl[\log \frac{q_{X|U}(x|u)}{q_X(x)} \\
&\qquad +\mu \left(\bar{\gamma} \log \frac{1}{q_{Y|U}(y|u)} + \gamma \log \frac{1}{q_{Z|U}(z|u)} \right) \Biggr], \\
&f^{(\alpha,\mu,\gamma,\lambda)}_{q_X||p_X,q_{XYZ|U}}(x,y,z|u) \\
&\triangleq \exp\left\{ -\lambda \omega^{(\alpha,\mu,\gamma)}_{q_X||p_X,q_{XYZ|U}}(x,y,z|u) \right\}, \\
&\Omega^{(\alpha,\mu,\gamma,\lambda)}(q|p_{XYZ}) \\
&\triangleq -\log \mathrm{E}_q \left[\exp\left\{ -\lambda \omega^{(\alpha,\mu,\gamma)}_{q_X||p_X,q_{XYZ|U}}(X,Y,Z|U) \right\} \right] \\
&= -\log \left[\sum_{u,x,y,z} q(u,x,y,z) f^{(\alpha,\mu,\gamma,\lambda)}_{q_X||p_X,q_{XYZ|U}}(x,y,z|u) \right], \\
&\Omega^{(\alpha,\mu,\gamma,\lambda)}(p_{XYZ}) \\
&\triangleq \min_{q \in \mathcal{Q}(p_{YZ|X}),} \Omega^{(\alpha,\mu,\gamma,\lambda)}(q|p_{XYZ}), \\
&F^{(\alpha,\mu,\gamma,\lambda)}(\alpha[R_1 + \mu(\bar{\gamma} R_2 + \gamma R_3)]|p_{XYZ}) \\
&\triangleq \frac{\Omega^{(\alpha,\mu,\gamma,\lambda)}(p_{XYZ}) - \alpha[R_1 + \mu(\bar{\gamma} R_2 + \gamma R_3)]}{1 + \lambda(1+\mu)}, \\
&F(R_1,R_2,R_3|p_{XYZ}) \\
&\triangleq \sup_{\substack{\lambda > 0, \\ (\alpha,\mu,\gamma) \\ \in [0,1]^3}} F^{(\alpha,\mu,\gamma,\lambda)}(\alpha[R_1 + \mu(\bar{\gamma} R_2 + \gamma R_3)]|p_{XYZ}), \\
&\overline{\mathcal{G}}(p_{XYZ}) \\
&\triangleq \left\{ (R_1,R_2,R_3,G) : G \geq F(R_1,R_2,R_3|p_{XYZ}) \right\}.
\end{aligned}$$

We can show that the above functions and sets satisfy the following property.

Property 7:

a) The cardinality bound $|\mathcal{U}| \leq |\mathcal{X}|$ in $\mathcal{Q}(p_{YZ|X})$ is sufficient to describe the quantity $\Omega^{(\alpha,\mu,\gamma,\lambda)}(p_{XYZ})$

b) Define a probability distribution $q^{(\lambda)} = q^{(\lambda)}_{UXYZ}$ by

$$\begin{aligned}
&q^{(\lambda)}(u,x,y,z) \\
&\triangleq \frac{q(u,x,y,z) \exp\left\{ -\lambda \omega^{(\alpha,\mu,\gamma)}_{q_X||p_X,q_{XYZ|U}}(x,y,z|u) \right\}}{\mathrm{E}_q \left[\exp\left\{ -\lambda \omega^{(\alpha,\mu,\gamma)}_{q_X||p_X,q_{XYZ|U}}(X,Y,Z|U) \right\} \right]}.
\end{aligned}$$

Then we have

$$\begin{aligned}
&\frac{\mathrm{d}}{\mathrm{d}\lambda} \Omega^{(\alpha,\mu,\gamma,\lambda)}(q|p_{XYZ}) \\
&\qquad = \mathrm{E}_{q^{(\lambda)}} \left[\omega^{(\alpha,\mu)}_{q_X||p_X,q_{XYZ|U}}(X,Y,Z|U) \right], \\
&\frac{\mathrm{d}^2}{\mathrm{d}\lambda^2} \Omega^{(\alpha,\mu,\gamma,\lambda)}(q|p_{XYZ}) \\
&\qquad = -\mathrm{Var}_{q^{(\lambda)}} \left[\omega^{(\alpha,\mu)}_{q_X||p_X,q_{XYZ|U}}(X,Y,Z|U) \right].
\end{aligned}$$

The second equality implies that $\Omega^{(\alpha,\mu,\gamma,\lambda)}(q|p_{XYZ})$ is a concave function of $\lambda > 0$.

c) Define

$$\begin{aligned}
&\rho = \rho(p_{XYZ}) \\
&\triangleq \max_{\substack{q \in \mathcal{Q}(p_{Y|X}), \\ (\alpha,\mu) \in [0,1]^2}} \mathrm{Var}_q \left[\omega^{(\alpha,\mu,\gamma)}_{q_X||p_X,q_{XYZ|U}}(X,Y,Z|U) \right].
\end{aligned}$$

Since

$$0 \leq \left[\omega^{(\alpha,\mu,\gamma)}_{q_X||p_X,q_{XYZ|U}}(x,y,z|u) \right]^2 < +\infty$$

for $(u,x,y,z) \in \mathcal{U} \times \mathcal{X} \times \mathcal{Y} \times \mathcal{Z}$, we have $\rho(p_{XYZ}) < \infty$. Then for any $\lambda \in (0,1]$, we have

$$\begin{aligned}
&\Omega^{(\alpha,\mu,\gamma,\lambda)}(q|p_{XYZ}) \\
&\geq \lambda \mathrm{E}_q \left[\omega^{(\alpha,\gamma,\mu)}_{q_X||p_X,q_{XYZ|U}}(X,Y,Z|U) \right] - \frac{1}{2} \rho(p_{XYZ}) \lambda^2.
\end{aligned}$$

Specifically, we have

$$\Omega^{(\alpha,\mu,\gamma,\lambda)}(p_{XYZ}) \geq \lambda \tilde{R}^{(\alpha,\mu)}(p_{XYZ}) - \frac{1}{2} \rho(p_{XYZ}) \lambda^2.$$

d) For any $\delta > 0$, there exists a positive number $\nu = \nu(\delta, |\mathcal{X}|, |\mathcal{Y}|, |\mathcal{Z}|) \in (0,1]$ such that for every positive $\tau \in (0, \nu]$, the condition $(R_1 + \tau, R_2 + \tau, R_3 + \tau) \notin \mathcal{R}(p_{XYZ})$ implies

$$F(R_1,R_2,R_3|p_{XYZ}) \geq \frac{\rho(p_{XYZ})}{2} \cdot g^2 \left(\frac{\tau^{3+\delta}}{2\rho(p_{XYZ})} \right) > 0,$$

Since the proof of Property 7 is quite similar to the proof of Property 3, we omit it.

Our main result is the following.

Theorem 6: For any $R_1, R_2, R_3 \geq 0$, any p_{XYZ}, and for any $(\varphi_1^{(n)}, \varphi_2^{(n)}, \varphi_3^{(n)}, \psi^{(n)}, \phi^{(n)})$ satisfying

$$\frac{1}{n} \log ||\varphi_i^{(n)}|| \leq R_i, i = 1,2,3,$$

we have

$$\begin{aligned}
&\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)}, \varphi_2^{(n)}, \varphi_3^{(n)}, \psi^{(n)}, \phi^{(n)}) \\
&\leq 7 \exp\left\{ -nF(R_1,R_2,R_3|p_{XYZ}) \right\}.
\end{aligned}$$

It follows from Theorem 6 and Property 7 part d) that if (R_1, R_2, R_3) is outside the capacity region, then the error probability of decoding goes to one exponentially and its exponent is not below $F(R_1,R_2,R_3|p_{XYZ})$. It immediately follows from Theorem 3 that we have the following corollary.

Corollary 4:

$$\begin{aligned}
&G(R_1,R_2,R_3|p_{XYZ}) \geq F(R_1,R_2,R_3|p_{XYZ}), \\
&\mathcal{G}(p_{XYZ}) \subseteq \overline{\mathcal{G}}(p_{XYZ}) \\
&= \left\{ (R_1,R_2,R_3,G) : G \geq F(R_1,R_2,R_3|p_{XYZ}) \right\}.
\end{aligned}$$

Proof of Theorem 6 will be given in the next section. The exponent function at rates outside the rate region was derived by Oohama and Han [7] for the separate source coding problem for correlated sources [6]. The techniques used by them is a method of types [8], which is not useful to prove Theorem 3. Some novel techniques based on the information spectrum method introduced by Han [9] are necessary to prove this theorem.

From Theorem 6 and Property 7 part d), we have the following corollary, which provides an explicit outer bound of $\mathcal{R}_{\mathrm{W}}(\varepsilon|p_{XYZ})$ with an asymptotically vanishing deviation from $\mathcal{R}_{\mathrm{W}}(p_{XYZ})=\mathcal{R}(p_{XYZ})$. The strong converse theorem immediately follows from this corollary.

From Theorem 6 and Property 7 part d) we have the following corollary.

Corollary 5: For each fixed $\varepsilon \in (0,1)$ and for any $\delta > 0$, there exists a positive integer n_0 with

$$n_0 = n_0(\varepsilon, \delta, |\mathcal{X}|, |\mathcal{Y}|, |\mathcal{Z}|, \rho(p_{XYZ}))$$

such that for $n \geq n_0$, we have

$$\mathcal{R}_{\mathrm{W}}(\varepsilon|p_{XYZ}) \subseteq \mathcal{R}(p_{XYZ}) - \tilde{\kappa}_n(1,1,1),$$

where

$$\tilde{\kappa}_n \stackrel{\triangle}{=} \left\{ \sqrt{\tfrac{8\rho(p_{XYZ})}{n} \log\left(\tfrac{7}{1-\varepsilon}\right)} + \tfrac{4}{n}\log\left(\tfrac{7}{1-\varepsilon}\right) \right\}^{\frac{1}{3+\delta}}.$$

It immediately follows from the above result that for any $\varepsilon \in (0,1)$, we have

$$\mathcal{R}_{\mathrm{W}}(\varepsilon|p_{XYZ}) = \mathcal{R}_{\mathrm{W}}(p_{XYZ}) = \mathcal{R}(p_{XYZ}).$$

APPENDIX

A. Properties of the Rate Regions

In this appendix we prove Property 1. Property 1 part a) can easily be proved by the definitions of the rate distortion regions. We omit the proofs of this part. In the following argument we prove part b).

Proof of Property 1 part b: We set

$$\underline{\mathcal{R}}_{\mathrm{AKW}}(m,\varepsilon|p_{XY}) = \bigcap_{n\geq m} \mathcal{R}_{\mathrm{AKW}}(n,\varepsilon|p_{XY}).$$

By the definitions of $\underline{\mathcal{R}}_{\mathrm{AKW}}(m,\varepsilon|p_{XY})$ and $\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})$, we have that $\underline{\mathcal{R}}_{\mathrm{AKW}}(m,\varepsilon|p_{XY}) \subseteq \mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})$ for $m \geq 1$. Hence we have that

$$\bigcup_{m\geq 1} \underline{\mathcal{R}}_{\mathrm{AKW}}(m,\varepsilon|p_{XY}) \subseteq \mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY}). \tag{40}$$

We next assume that $(R_1,R_2) \in \mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})$. Set

$$\begin{aligned}&\mathcal{R}^{(\delta)}_{\mathrm{AKW}}(\varepsilon|p_{XY})\\ &\stackrel{\triangle}{=} \{(R_1+\delta, R_2+\delta) : (R_1,R_2) \in \mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})\}\end{aligned}$$

Then, by the definitions of $\mathcal{R}_{\mathrm{AKW}}(n,\varepsilon\ |p_{XY})$ and $\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})$, we have that for any $\delta > 0$, there exists $n_0(\varepsilon,\delta)$ such that for any $n \geq n_0(\varepsilon,\delta)$,

$$(R+\delta, R_2+\delta) \in \mathcal{R}_{\mathrm{AKW}}(n,\varepsilon|p_{XY}),$$

which implies that

$$\begin{aligned}\mathcal{R}^{(\delta)}_{\mathrm{AKW}}(\varepsilon|p_{XY}) &\subseteq \bigcup_{n\geq n_0(\varepsilon,\delta)} \mathcal{R}_{\mathrm{AKW}}(n,\varepsilon|p_{XY})\\ &= \underline{\mathcal{R}}_{\mathrm{AKW}}(n_0(\delta),\varepsilon|p_{XY})\\ &\subseteq \mathrm{cl}\left(\bigcup_{m\geq 1} \underline{\mathcal{R}}_{\mathrm{AKW}}(m,\varepsilon|p_{XY})\right).\end{aligned} \tag{41}$$

Here we assume that there exists a pair (R_1,R_2) belonging to $\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})$ such that

$$(R_1,R_2) \notin \mathrm{cl}\left(\bigcup_{m\geq 1} \underline{\mathcal{R}}_{\mathrm{AKW}}(m,\varepsilon|p_{XY})\right). \tag{42}$$

Since the set in the right hand side of (42) is a closed set, we have

$$(R_1+\delta, R_2+\delta) \notin \mathrm{cl}\left(\bigcup_{m\geq 1} \underline{\mathcal{R}}_{\mathrm{AKW}}(m,\varepsilon|p_{XY})\right) \tag{43}$$

for some small $\delta > 0$. On the other hand we have $(R+\delta, R_2+\delta) \in \mathcal{R}^{(\delta)}_{\mathrm{AKW}}(\varepsilon|p_{XY})$, which contradicts (41). Thus we have

$$\begin{aligned}&\bigcup_{m\geq 1} \underline{\mathcal{R}}_{\mathrm{AKW}}(m,\varepsilon|p_{XY})\\ &\subseteq \mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY}) \subseteq \mathrm{cl}\left(\bigcup_{m\geq 1} \underline{\mathcal{R}}_{\mathrm{AKW}}(m,\varepsilon|p_{XY})\right).\end{aligned} \tag{44}$$

Note here that $\mathcal{R}_{\mathrm{AKW}}(\varepsilon|p_{XY})$ is a closed set. Then from (44), we conclude that

$$\begin{aligned}\mathcal{R}_{\mathrm{AKW}}(\varepsilon|W) &= \mathrm{cl}\left(\bigcup_{m\geq 1} \underline{\mathcal{R}}_{\mathrm{AKW}}(m,\varepsilon|p_{XY})\right)\\ &= \mathrm{cl}\left(\bigcup_{m\geq 1}\bigcap_{n\geq m} \mathcal{R}_{\mathrm{AKW}}(n,\varepsilon|p_{XY})\right),\end{aligned}$$

completing the proof. ∎

B. Cardinality Bound on Auxiliary Random Variables

We first prove the following lemma.

Lemma 9:

$$\begin{aligned}&\underline{R}^{(\mu)}(p_{XY}) \stackrel{\triangle}{=} \min_{p\in\mathcal{P}(p_{XY})} \{\bar{\mu} I_p(X;U) + \mu H_p(Y|U)\}\\ &= R^{(\mu)}(p_{XY}) \stackrel{\triangle}{=} \min_{p\in\mathcal{P}_{\mathrm{sh}}(p_{XY})} \{\bar{\mu} I_p(X;U) + \mu H_p(Y|U)\}.\end{aligned}$$

Proof: We bound the cardinality $|\mathcal{U}|$ of U to show that the bound $|\mathcal{U}| \leq |\mathcal{X}|$ is sufficient to describe $\underline{R}^{(\mu)}(p_{XY})$. Observe that

$$p_X(x) = \sum_{u\in\mathcal{U}} p_U(u) p_{X|U}(x|u), \tag{45}$$

$$\bar{\mu} I_p(X;U) + \mu H_p(Y|U) = \sum_{u\in\mathcal{U}} p_U(u)\pi(p_{X|U}(\cdot|u)), \tag{46}$$

where

$$\begin{aligned}\pi(p_{X|U}(\cdot|u)) &\stackrel{\triangle}{=} \sum_{(x,y)\in\mathcal{X}\times\mathcal{Y}} p_{X|U}(x|u)p_{Y|X}(y|x)\\ &\times \log\left\{\frac{p^{\bar{\mu}}_{X|U}(x|u)}{p^{\bar{\mu}}_X(x)}\left[\sum_{\tilde{x}\in\mathcal{X}} p_{Y|X}(y|\tilde{x})p_{X|U}(\tilde{x}|u)\right]^{-\mu}\right\}.\end{aligned}$$

For each $u \in \mathcal{U}$, $\pi(p_{X|U}(\cdot|u))$ is a continuous function of $p_{X|U}(\cdot|u)$. Then by the support lemma,

$$|\mathcal{U}| \leq |\mathcal{X}| - 1 + 1 = |\mathcal{X}|$$

is sufficient to express $|\mathcal{X}|-1$ values of (45) and one value of (46). ■

Next we prove the following lemma.

Lemma 10: For each integer $n \geq 2$, we have

$$\begin{aligned}
&\hat{\Omega}_n^{(\alpha,\mu,\lambda)}(p_{XY}) \\
&\triangleq \min_{\substack{q=q_{UXY}: U \leftrightarrow X \leftrightarrow Y, \\ q_{Y|X}=p_{Y|X}, \\ |\mathcal{U}| \leq |\mathcal{M}_1||\mathcal{X}|^{n-1}}} \Omega^{(\alpha,\mu,\lambda)}(q|p_{XY}) \\
&= \min_{\substack{q=q_{UXY}: U \leftrightarrow X \leftrightarrow Y, \\ q_{Y|X}=p_{Y|X}, \\ |\mathcal{U}| \leq |\mathcal{X}|}} \Omega^{(\alpha,\mu,\lambda)}(q|p_{XY}) \\
&\triangleq \Omega^{(\alpha,\mu,\lambda)}(p_{XY}).
\end{aligned}$$

Proof: We bound the cardinality $|\mathcal{U}|$ of U to show that the bound $|\mathcal{U}| \leq |\mathcal{X}|$ is sufficient to describe $\hat{\Omega}_n^{(\alpha,\mu,\lambda)}(p_{XY})$. Observe that

$$q_X(x) = \sum_{u \in \mathcal{U}} q_U(u) q_{X|U}(x|u), \tag{47}$$

$$\begin{aligned}
&\exp\left\{-\Omega^{(\alpha,\mu,\lambda)}(q|p_{XY})\right\} \\
&\quad = \sum_{u \in \mathcal{U}} q_U(u) \Pi(q_{X|U}(\cdot|u)),
\end{aligned} \tag{48}$$

where

$$\begin{aligned}
\Pi(q_{X|U}(\cdot|u)) &\triangleq \sum_{(x,y) \in \mathcal{X} \times \mathcal{Y}} q_{X|U}^{1-\lambda}(x|u) q_{Y|X}(y|x) \\
&\quad \times \exp\left\{-\omega_{q_X||p_X, q_{XY|U}}^{(\alpha,\mu)}(x,y|u)\right\}.
\end{aligned}$$

For each $u \in \mathcal{U}$, $\Pi(q_{X|U}(\cdot|u))$ is a continuous function of $q_{X|U}(\cdot|u)$. Then by the support lemma,

$$|\mathcal{U}| \leq |\mathcal{X}| - 1 + 1 = |\mathcal{X}|$$

is sufficient to express $|\mathcal{X}|-1$ values of (47) and one value of (48). ■

C. *Supporting Hyperplain Expressions of* $\mathcal{R}(p_{XY})$

In this appendix we prove Property 2 part b). We first prepare a lemma useful to prove this property.

From the convex property of the region $\mathcal{R}(p_{XY})$, we have the following lemma.

Lemma 11: Suppose that $(\hat{R}_1, \hat{R}_2)$ does not belong to $\mathcal{R}(p_{XY})$. Then there exist $\epsilon, \mu^* > 0$ such that for any $(R_1, R_2) \in \mathcal{R}(p_{XY})$ we have

$$\bar{\mu}(R_1 - \hat{R}_1) + \mu^*(R_2 - \hat{R}_2) - \epsilon \geq 0.$$

Proof of this lemma is omitted here. Lemma 11 is equivalent to the fact that if the region $\mathcal{R}(p_{XY})$ is a convex set, then for any point $(\hat{R}_1, \hat{R}_2)$ outside the region $\mathcal{R}(p_{XY})$, there exists a line which separates the point $(\hat{R}_1, \hat{R}_2)$ from the region $\mathcal{R}(p_{XY})$. Lemma 11 will be used to prove Property 2 part b).

Proof of Property 2 part b): We first recall the following definitions of $\mathcal{P}^*(p_{XY})$ and $\mathcal{P}(p_{XY})$:

$$\begin{aligned}
\mathcal{P}(p_{XY}) &\triangleq \{p_{UXY} : |\mathcal{U}| \leq |\mathcal{X}| + 1, U \leftrightarrow X \leftrightarrow Y\}, \\
\mathcal{P}_{\mathrm{sh}}(p_{XY}) &\triangleq \{p_{UXY} : |\mathcal{U}| \leq |\mathcal{X}|, U \leftrightarrow X \leftrightarrow Y\}.
\end{aligned}$$

We prove $\mathcal{R}_{\mathrm{sh}}(p_{XY}) \subseteq \mathcal{R}(p_{XY})$. We assume that $(\hat{R}_1, \hat{R}_2) \notin \mathcal{R}(p_{XY})$. Then by Lemma 11, there exist $\epsilon > 0$ and $\mu^* \in [0,1]$ such that for any $(R_1, R_2) \in \mathcal{R}(p_{XY})$, we have

$$\bar{\mu}^* \hat{R}_1 + \mu^* \hat{R}_2 \leq \bar{\mu}^* R_1 + \mu^* R_2 - \epsilon.$$

Hence we have

$$\begin{aligned}
\bar{\mu}^* \hat{R}_1 + \mu^* \hat{R}_2 &\leq \min_{(R_1,R_2) \in \mathcal{R}(p_{XY})} \{\bar{\mu}^* R_1 + \mu^* R_2\} - \epsilon \\
&\overset{\text{(a)}}{=} \min_{p \in \mathcal{P}(p_{XY})} \{\bar{\mu}^* I_p(U;X|Y) + \mu^* H_p(Y|U)\} - \epsilon \\
&\leq \min_{p \in \mathcal{P}_{\mathrm{sh}}(p_{XY})} \{\bar{\mu}^* I_p(U;X) + \mu^* H_p(Y|U)\} - \epsilon \\
&= R^{(\mu^*)}(p_{XY}) - \epsilon.
\end{aligned} \tag{49}$$

Step (a) follows from the definition of $\mathcal{R}(p_{XY})$. The inequality (49) implies that $(\hat{R}_1, \hat{R}_2) \notin \mathcal{R}_{\mathrm{sh}}(p_{XY})$. Thus $\mathcal{R}_{\mathrm{sh}}(p_{XY}) \subseteq \mathcal{R}(p_{XY})$ is concluded. We next prove $\mathcal{R}(\ p_{XY}) \subseteq \mathcal{R}_{\mathrm{sh}}(p_{XY})$. We assume that $(R_1,\ R_2) \in \mathcal{R}(p_{XY})$. Then there exists $q \in \mathcal{P}\ (p_{XY})$ such that

$$R_1 \geq I_q(X;U), R_2 \geq H_q(Y|U). \tag{50}$$

Then, for each $\mu > 0$ and for $(R_1, R_2) \in \mathcal{R}(p_{XY})$, we have the following chain of inequalities:

$$\begin{aligned}
\bar{\mu}R_1 + \mu R_2 &\overset{\text{(a)}}{\geq} \bar{\mu} I_q(X;U) + \mu H_q(Y|U) \\
&\geq \min_{q \in \mathcal{P}(p_{XY})} [\bar{\mu} I_q(X;U) + \mu H_q(Y|U)] \\
&= R^{(\mu)}(p_{XY}).
\end{aligned}$$

Step (a) follows from (50). Hence we have $\mathcal{R}(p_{XY}) \subseteq \mathcal{R}_{\mathrm{sh}}(p_{XY})$. We next prove the second inequality of (6) in Property 2. We have the following chain of inequalities.

$$\begin{aligned}
&R^{(\mu)}(p_{XY}). \\
&\geq \min_{q \in \mathcal{P}_{\mathrm{sh}}(p_{XY})} [\bar{\mu} I_q(X;U) + \mu H_q(Y|U)] \\
&\overset{\text{(a)}}{=} \min_{q \in \mathcal{P}(p_{XY})} \left\{ \frac{(1+\mu)\bar{\alpha}}{\alpha} D(q_X||p_X) \right. \\
&\qquad \left. + \bar{\mu} I_q(X;U) + \mu H_q(Y|U) \right\} \\
&\geq \min_{q \in \mathcal{Q}(p_{Y|X})} \left\{ \frac{(1+\mu)\bar{\alpha}}{\alpha} D(q_X||p_X) \right. \\
&\qquad \left. + \bar{\mu} I_q(X;U) + \mu H_q(Y|U) \right\} \\
&= \frac{1}{\alpha} \tilde{R}^{(\alpha,\mu)}(p_{XY}).
\end{aligned}$$

Step (a) follows from that when $q \in \mathcal{P}_{\mathrm{sh}}(p_{XY})$, we have $D(q_X||p_X) = 0$. We finally prove the first inequality of (6) in Property 2. Let

$$q_{\alpha,\mu}^* = q_{UXY,\alpha,\mu}^* \in \mathcal{Q}(p_{Y|X})$$

be the probability distribution attaing the minimum in the definition of $\tilde{R}^{(\alpha,\mu)}(p_{XY})$. Let $\hat{q}_{\alpha,\mu} = \hat{q}_{UXYZ,\alpha,\mu}$ be a probability disitribution with the form

$$\hat{q}_{UXY,\alpha,\mu}(u,x,y) = q_{U|X,\alpha,\mu}^*(u|x) p_X(x) p_{Y|X}(y|x).$$

By definition, we have $\hat{q}_{\alpha,\mu} \in \mathcal{P}_{\rm sh}(p_{XY})$. Then we have the following chain of inequalities.

$$\begin{aligned}
&\bar{\alpha}D(q^*_{\alpha,\mu}||\hat{q}_{\alpha,\mu})\\
&= \bar{\alpha}D(q^*_{X,\alpha,\mu}||p_X)\\
&\le (1+\mu)\bar{\alpha}D(q^*_{X,\alpha,\mu}||p_X)\\
&\quad +\alpha[\bar{\mu}I_{q^*_{\alpha,\mu}}(X;U)+\mu H_{q^*_{\alpha,\mu}}(Y|U)]\\
&= \tilde{R}^{(\alpha,\mu)}(p_{XY})\\
&\le (1+\mu)\bar{\alpha}[D(\hat{q}_{X,\alpha,\mu}||p_X)\\
&\quad +\alpha[\bar{\mu}I_{\hat{q}_{\alpha,\mu}}(X;U)+\mu H_{\hat{q}_{\alpha,\mu}}(Y|U)]\\
&= \alpha[\bar{\mu}I_{\hat{q}_{\alpha,\mu}}(X;U)+\mu H_{\hat{q}_{\alpha,\mu}}(Y|U)]\\
&\le \alpha\log(|\mathcal{X}||\mathcal{Y}|).
\end{aligned} \tag{51}$$

For simplicity of notation we set $\xi \overset{\triangle}{=} \log(|\mathcal{X}||\mathcal{Y}|)$. From (51) we have

$$D(q^*_{\alpha,\mu}||\hat{q}_{\alpha,\mu}) \le \xi\frac{\alpha}{\bar{\alpha}}. \tag{52}$$

By the Pinsker's inequality we have

$$\frac{1}{2}(||q^*_{\alpha,\mu}-\hat{q}_{\alpha,\mu}||_1)^2 \le D(q^*_{\alpha,\mu}||\hat{q}_{\alpha,\mu}) \tag{53}$$

From (52) and (53), we obtain

$$||q^*_{\alpha,\mu}-\hat{q}_{\alpha,\mu}||_1 \le \sqrt{2\xi\frac{\alpha}{\bar{\alpha}}} \le \frac{1}{2}. \tag{54}$$

for any $\alpha \in (0,\alpha_0]$ with

$$\alpha_0 = (8\xi+1)^{-1} = [8\log(|\mathcal{X}||\mathcal{Y}|)+1]^{-1}. \tag{55}$$

The bound (54) implies that for any $A \subseteq \{U,X,Y,Z\}$, we have

$$||q^*_{A,\alpha,\mu}-\hat{q}_{A,\alpha,\mu}||_1 \le \sqrt{2\xi\frac{\alpha}{\bar{\alpha}}} \le \frac{1}{2}. \tag{56}$$

Then for any $\alpha \in (0,\alpha_0]$, we have the following two chains of inequalities:

$$\begin{aligned}
&|I_{q^*_{\alpha,\mu}}(U;X)-I_{\hat{q}_{\alpha,\mu}}(U;X)|\\
&\le |H_{q^*_{\alpha,\mu}}(U)-H_{\hat{q}_{\alpha,\mu}}(U)|+|H_{q^*_{\alpha,\mu}}(X)-H_{\hat{q}_{\alpha,\mu}}(X)|\\
&\quad +|H_{q^*_{\alpha,\mu}}(UX)-H_{\hat{q}_{\alpha,\mu}}(UX)|\\
&\overset{\rm (a)}{\le} -\sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\left[\log\left(\sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\cdot\frac{1}{|\mathcal{U}|}\right)+\log\left(\sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\cdot\frac{1}{|\mathcal{X}|}\right)\right.\\
&\qquad \left.+\log\left(\sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\cdot\frac{1}{|\mathcal{U}||\mathcal{X}|}\right)\right]\\
&= \sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\log\left\{\left(\frac{\bar{\alpha}}{2\xi\alpha}\right)^{\frac{3}{2}}(|\mathcal{U}||\mathcal{X}|)^2\right\},
\end{aligned} \tag{57}$$

$$\begin{aligned}
&|H_{q^*_{\alpha,\mu}}(Y|U)-H_{\hat{q}_{\alpha,\mu}}(Y|U)|\\
&\le |H_{q^*_{\alpha,\mu}}(YU)-H_{\hat{q}_{\alpha,\mu}}(YU)|\\
&\quad +|H_{q^*_{\alpha,\mu}}(U)-H_{\hat{q}_{\alpha,\mu}}(U)|\\
&\overset{\rm (b)}{\le} -\sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\left[\log\left(\sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\cdot\frac{1}{|\mathcal{U}||\mathcal{Y}|}\right)\right.\\
&\qquad \left.+\log\left(\sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\cdot\frac{1}{|\mathcal{U}|}\right)\right]\\
&\le \sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\log\left\{\left(\frac{\bar{\alpha}}{2\xi\alpha}\right)|\mathcal{U}|^2|\mathcal{Y}|\right\}.
\end{aligned} \tag{58}$$

Steps (a) and (b) follow from (56) and LEMMA 2.7 in Section 1.2 in Csiszár and Körner [8]. Combining (57) and (58), we have

$$\begin{aligned}
&\Bigl|\bar{\mu}I_{q^*_{\alpha,\mu}}(X;U)+\mu H_{q^*_{\alpha,\mu}}(Y|U)\\
&\quad -\left(\bar{\mu}I_{\hat{q}_{\alpha,\mu}}(X;U)+\mu H_{\hat{q}_{\alpha,\mu}}(Y|U)\right)\Bigr|\\
&\le \bar{\mu}|I_{q^*_{\alpha,\mu}}(X;U)-I_{\hat{q}_{\alpha,\mu}}(X;U)|\\
&\quad +\mu|H_{q^*_{\alpha,\mu}}(Y|U)-H_{\hat{q}_{\alpha,\mu}}(Y|U)|\\
&\le \sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\log\left\{\left(\frac{\bar{\alpha}}{2\xi\alpha}\right)^{\frac{3-\mu}{2}}|\mathcal{U}|^2|\mathcal{X}|^{2\bar{\mu}}|\mathcal{Y}|^{\mu}\right\}\\
&\le \sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\log\left\{\left(\frac{\bar{\alpha}}{2\xi\alpha}\right)^{\frac{3}{2}}|\mathcal{U}|^2|\mathcal{X}|^2|\mathcal{Y}|\right\}\\
&\overset{\rm (a)}{\le} \sqrt{2\xi\frac{\alpha}{\bar{\alpha}}}\log\left\{\left(\frac{\bar{\alpha}}{2\xi\alpha}\right)^{\frac{3}{2}}|\mathcal{X}|^4|\mathcal{Y}|\right\}\\
&= c_1\sqrt{\frac{\alpha}{\bar{\alpha}}}\log\left(c_2\frac{\bar{\alpha}}{\alpha}\right).
\end{aligned} \tag{59}$$

Step (a) follows from the cardinality bound $|\mathcal{U}| \le |\mathcal{X}|$ we have for $q^*_{\alpha,\mu} \in \mathcal{Q}(p_{Y|X})$. Then we have

$$\begin{aligned}
&\frac{1}{\alpha}\tilde{R}^{(\alpha,\mu)}(p_{XY})\\
&= (1+\mu)\frac{\bar{\alpha}}{\alpha}D(q^*_{X,\alpha,\mu}||p_X)\\
&\quad +I_{q^*_{\alpha,\mu}}(X;U)+\mu H_{q^*_{\alpha,\mu}}(Y|U)\\
&\ge I_{q^*_{\alpha,\mu}}(X;U)+\mu H_{q^*_{\alpha,\mu}}(Y|U)\\
&\overset{\rm (a)}{\ge} I_{\hat{q}_{\alpha,\mu}}(X;U)+\mu H_{\hat{q}_{\alpha,\mu}}(Y|U)-c_1\sqrt{\frac{\alpha}{\bar{\alpha}}}\log\left(c_2\frac{\bar{\alpha}}{\alpha}\right)\\
&\overset{\rm (b)}{\ge} \min_{q\in\mathcal{P}_{\rm sh}(p_{XY})}\{I_q(X;U)+\mu H_q(Y|U)\}\\
&\quad -c_1\sqrt{\frac{\alpha}{\bar{\alpha}}}\log\left(c_2\frac{\bar{\alpha}}{\alpha}\right)\\
&= R^{(\mu)}(p_{XY})-c_1\sqrt{\frac{\alpha}{\bar{\alpha}}}\log\left(c_2\frac{\bar{\alpha}}{\alpha}\right).
\end{aligned} \tag{60}$$

Step (a) follows from (59). Step (b) follows from that $\hat{q}_{\alpha,\mu} \in \mathcal{Q}(p_{Y|X})$. ■

D. *Proof of Property 3 parts b), c), and d)*

In this appendix we prove Property 3 b),c), and d).

Proof of Property 3b),c), and d): We first prove part b). For simplicity of notations, set

$$\begin{aligned}
&\underline{a} \overset{\triangle}{=} (u,x,y), \underline{A} \overset{\triangle}{=} (U,X,Y), \underline{\mathcal{A}} \overset{\triangle}{=} \mathcal{U}\times\mathcal{X}\times\mathcal{Y},\\
&\omega^{(\mu)}_{q_{XY|U}}(x,y|u) \overset{\triangle}{=} g(\underline{a}),\\
&\Omega^{(\alpha,\mu,\lambda)}_q(XY|U) \overset{\triangle}{=} \xi(\lambda).
\end{aligned}$$

Then we have

$$\xi(\lambda) = -\log\left[\sum_{\underline{a}\in\underline{\mathcal{A}}} p_{\underline{A}}(\underline{a}){\rm e}^{-\lambda g(\underline{a})}\right].$$

We set

$$q^{(\lambda)}(u,x,y,z) \overset{\triangle}{=} p^{(\lambda)}_{\underline{A}}(\underline{a}).$$

Then $p_{\underline{A}}^{(\lambda)}(\underline{a}), \underline{a} \in \mathcal{A}$ has the following form:

$$p_{\underline{A}}^{(\lambda)}(\underline{a}) = \mathrm{e}^{\xi(\lambda)} p_{\underline{A}}(\underline{a}) \mathrm{e}^{-\lambda g(\underline{a})}.$$

By simple computations we have

$$\begin{aligned} \xi'(\lambda) &= \mathrm{e}^{\xi(\lambda)} \left[\sum_{\underline{a} \in \underline{\mathcal{A}}} p_{\underline{A}}(\underline{a}) g(\underline{a}) \mathrm{e}^{-\lambda g(\underline{a})} \right], \\ &= \sum_{\underline{a} \in \underline{\mathcal{A}}} p_{\underline{A}}^{(\lambda)}(\underline{a}) g(\underline{a}). \end{aligned}$$

$$\begin{aligned} &\xi''(\lambda) = -\mathrm{e}^{2\xi(\lambda)} \\ &\times \left[\sum_{\underline{a}, \underline{b} \in \underline{\mathcal{A}}} p_{\underline{A}}(\underline{a}) p_{\underline{A}}(\underline{b}) \frac{\{g(\underline{a}) - g(\underline{b})\}^2}{2} \mathrm{e}^{-\lambda\{g(\underline{a}) + g(\underline{b})\}} \right] \\ &= - \sum_{\underline{a}, \underline{b} \in \underline{\mathcal{A}}} p_{\underline{A}}^{(\lambda)}(\underline{a}) p_{\underline{A}}^{(\lambda)}(\underline{b}) \frac{\{g(\underline{a}) - g(\underline{b})\}^2}{2} \\ &= - \sum_{\underline{a} \in \underline{\mathcal{A}}} p_{\underline{A}}^{(\lambda)}(\underline{a}) g^2(\underline{a}) + \left[\sum_{\underline{a} \in \underline{\mathcal{A}}} p_{\underline{A}}^{(\lambda)}(\underline{a}) g(\underline{a}) \right]^2. \end{aligned}$$

We next prove the part c). Let $\lambda \in [0, 1]$. By the Taylor expansion of $\Omega^{(\alpha,\mu,\lambda)}(q|p_{XY})$ with respect to λ around $\lambda = 0$, we have

$$\begin{aligned} &\Omega^{(\alpha,\mu,\lambda)}(q|p_{XY}) \\ &= \xi(\lambda) = \xi(0) + \xi'(0)\lambda + \frac{1}{2}\xi''(0)(\tau\lambda)^2 \\ &= \mathrm{E}_q \left[\omega^{(\alpha,\mu)}_{q_X || p_X, q_{XY|U}}(X, Y|U) \right] \\ &\quad -\frac{1}{2}(\tau\lambda)^2 \mathrm{Var}_q \left[\omega^{(\alpha,\mu)}_{q_X || p_X, q_{XY|U}}(X, Y|U) \right] \end{aligned}$$

for some $\tau \in [0, 1]$. Then by the definition of ρ, we have

$$\begin{aligned} &\Omega^{(\alpha,\mu,\lambda)}(q|p_{XY}) \\ &\geq \lambda \mathrm{E}_q \left[\omega^{(\alpha,\mu)}_{q_X || p_X, q_{XY|U}}(X, Y|U) \right] - \frac{1}{2}\rho\lambda^2. \end{aligned}$$

The second inequality is obvious from the first inequality. We finally prove the part d). By the hyperplane expression $\mathcal{R}_{\mathrm{sh}}(p_{XY})$ of $\mathcal{R}(p_{XY})$ stated Property 2 part b) we have that when $(R_1 + \tau, R_2 + \tau) \notin \mathcal{R}(p_{XY})$, we have

$$\bar{\mu}^*(R_1 + \tau) + \mu^*(R_2 + \tau) < R^{(\mu^*)}(p_{XY}) \tag{61}$$

for some $\mu^* \in [0, 1]$. Let $\alpha_0 = \alpha_0(|\mathcal{X}|, |\mathcal{Y}|)$ be the quantity defined by (7) in Property 2 part b). Then from (61), we have

$$\begin{aligned} &\bar{\mu}^* R_1 + \mu^* R_2 + \tau \leq R^{(\mu^*)}(p_{XY}) \\ &\overset{\text{(a)}}{\leq} \frac{1}{\alpha} \tilde{R}^{(\alpha,\mu^*)}(p_{XY}) + c_1 \sqrt{\frac{\alpha}{\bar{\alpha}}} \log\left(c_2 \frac{\bar{\alpha}}{\alpha} \right). \end{aligned} \tag{62}$$

Step (a) follows from the first inequality of (6) in Property 2 part b). Fix any $\delta > 0$. We choose $\alpha = \tau^{2+\delta}$. For this choice of α, we have (62) for any $\tau \in (0, \alpha_0^{\frac{1}{2+\delta}}]$ and

$$\begin{aligned} &c_1 \sqrt{\frac{\alpha}{\bar{\alpha}}} \log\left(c_2 \frac{\bar{\alpha}}{\alpha} \right) \times \frac{1}{\tau} \\ &= c_1 \frac{\tau^{\frac{\delta}{2}}}{\sqrt{1 - \tau^{2+\delta}}} \log\left(c_2 \frac{1 - \tau^{2+\delta}}{\tau^{2+\delta}} \right) \to 0 \text{ as } \tau \to 0. \end{aligned}$$

Hence there exists a positive $\nu = \nu(\delta, |\mathcal{X}|, |\mathcal{Y}|)$ with $\nu \leq \alpha_0^{\frac{1}{2+\delta}}$ such that for $\tau \in (0, \nu] \subseteq (0, \alpha_0^{\frac{1}{2+\delta}}]$,

$$c_1 \cdot \sqrt{\frac{\alpha}{\bar{\alpha}}} \log\left(c_2 \frac{\bar{\alpha}}{\alpha} \right) \leq \frac{\tau}{2}. \tag{63}$$

The above inequality together with (62) yields that for $\tau \in (0, \nu]$,

$$\bar{\mu}^* R_1 + \mu^* R_2 + \frac{1}{2}\tau \leq \frac{1}{\tau^{2+\delta}} \tilde{R}^{(\tau^{2+\delta},\mu)}(p_{XY}). \tag{64}$$

Then for each positive τ with $\tau \in (0, \nu]$, we have the following chain of inequalities:

$$\begin{aligned} &F(R_1, R_2|p_{XY}) \\ &\geq \sup_{\lambda > 0} F^{(\tau^{2+\delta},\mu^*,\lambda)}(\bar{\mu}^* R_1 + \mu^* R_2 | p_{XY}) \\ &= \sup_{\lambda > 0} \frac{\Omega^{(\tau^{2+\delta},\mu^*,\lambda)}(p_{XY}) - \lambda\tau^{2+\delta}(\bar{\mu}^* R_1 + \mu^* R_2)}{1 + \lambda(1 + \mu^*)} \\ &\overset{\text{(a)}}{\geq} \sup_{\lambda > 0} \frac{1}{1 + 2\lambda} \left\{ -\frac{1}{2}\rho\lambda^2 + \lambda \tilde{R}^{(\tau^{2+\delta},\mu^*)}(p_{XY}) \right. \\ &\qquad\qquad \left. -\lambda\tau^{2+\delta}(\bar{\mu}^* R_1 + \mu^* R_2) \right\} \\ &\overset{\text{(b)}}{\geq} \sup_{\lambda > 0} \frac{1}{1 + 2\lambda} \left\{ -\frac{1}{2}\rho\lambda^2 + \frac{1}{2}\tau^{3+\delta}\lambda \right\} \overset{\text{(c)}}{=} \frac{\rho}{2} g^2 \left(\frac{\tau^{3+\delta}}{2\rho} \right). \end{aligned}$$

Step (a) follows from Property 3 part b). Step (b) follows from (64). Step (c) follows from an elementary computation. This completes the proof of Property 3 part d). ■

E. *Proof of Lemma 1*

To prove Lemma 1, we prepare a lemma. Set

$$\mathcal{A}_n \triangleq \left\{ (s, x^n, y^n) : \frac{1}{n} \log \frac{p_{V^n}(s, x^n, y^n)}{q_{V^n}(s, x^n, y^n)} \geq -\eta \right\}.$$

Furthermore, set

$$\begin{aligned} &\tilde{\mathcal{B}}_n \triangleq \left\{ x^n : \frac{1}{n} \log \frac{p_{X^n}(x^n)}{q_{X^n}(x^n)} \geq -\eta \right\}, \\ &\mathcal{B}_n \triangleq \tilde{\mathcal{B}}_n \times \mathcal{M}_1 \times \mathcal{Y}^n, \mathcal{B}_n^{\mathrm{c}} \triangleq \tilde{\mathcal{B}}_n^{\mathrm{c}} \times \mathcal{M}_1 \times \mathcal{Y}^n, \\ &\tilde{\mathcal{C}}_n \triangleq \{ (s, x^n) : s = \varphi_1^{(n)}(x^n), \\ &\qquad\qquad p_{X^n|S}(x^n|s) \leq M_1 \mathrm{e}^{n\eta} p_{X^n}(x^n) \}, \\ &\mathcal{C}_n \triangleq \tilde{\mathcal{C}}_n \times \mathcal{Y}^n, \mathcal{C}_n^{\mathrm{c}} \triangleq \tilde{\mathcal{C}}_n^{\mathrm{c}} \times \mathcal{Y}^n, \\ &\mathcal{D}_n \triangleq \{ (s, x^n, y^n) : s = \varphi_1^{(n)}(x^n), \\ &\qquad\qquad p_{Y^n|S}(y^n|s) \geq (1/M_2) \mathrm{e}^{-n\eta} \}, \\ &\mathcal{E}_n \triangleq \{ (s, x^n, y^n) : s = \varphi_1^{(n)}(x^n), \\ &\qquad\qquad \psi^{(n)}(\varphi_1^{(n)}(x^n), \varphi_2^{(n)}(y^n)) = y^n \}. \end{aligned}$$

Then we have the following lemma.

Lemma 12:

$$\begin{aligned} &p_{SX^nY^n}(\mathcal{A}_n^{\mathrm{c}}) \leq \mathrm{e}^{-n\eta}, p_{SX^nY^n}(\mathcal{B}_n^{\mathrm{c}}) \leq \mathrm{e}^{-n\eta}, \\ &p_{SX^nY^n}(\mathcal{C}_n^{\mathrm{c}}) \leq \mathrm{e}^{-n\eta}, p_{SX^nY^n}(\mathcal{D}_n^{\mathrm{c}} \cap \mathcal{E}_n) \leq \mathrm{e}^{-n\eta}. \end{aligned}$$

Proof: We first prove the first inequality.

$$
\begin{aligned}
p_{SX^nY^n}(\mathcal{A}_n^{\mathrm{c}}) &= \sum_{(s,x^n,y^n)\in\mathcal{A}_n^{\mathrm{c}}} p_{V_n}(s,x^n,y^n) \\
&\overset{\text{(a)}}{\leq} \sum_{(s,x^n,y^n)\in\mathcal{A}_n^{\mathrm{c}}} \mathrm{e}^{-n\eta} q_{V_n}(s,x^n,y^n) \\
&= \mathrm{e}^{-n\eta} q_{V_n}(\mathcal{A}_n^{\mathrm{c}}) \leq \mathrm{e}^{-n\eta}.
\end{aligned}
$$

Step (a) follows from the definition of $\mathcal{A}_n$. On the second iniquality we have

$$
\begin{aligned}
p_{SX^nY^n}(\mathcal{B}_n^{\mathrm{c}}) &= p_{X^n}(\tilde{\mathcal{B}}_n^{\mathrm{c}}) = \sum_{x^n\in\tilde{\mathcal{B}}_n^{\mathrm{c}}} p_{X_n}(x^n) \\
&\overset{\text{(a)}}{\leq} \sum_{x^n\in\tilde{\mathcal{B}}_n^{\mathrm{c}}} \mathrm{e}^{-n\eta} q_{X^n}(x^n) \\
&= \mathrm{e}^{-n\eta} q_{X^n}\left(\tilde{\mathcal{B}}_n^{\mathrm{c}}\right) \leq \mathrm{e}^{-n\eta}.
\end{aligned}
$$

Step (a) follows from the definition of $\mathcal{B}_n$. We next prove the third inequality.

$$
\begin{aligned}
&p_{SX^nY^n}(\mathcal{C}_n^{\mathrm{c}}) = p_{SX^n}(\tilde{\mathcal{C}}_n^{\mathrm{c}}) \\
&= \sum_{s\in\mathcal{M}_1} \sum_{\substack{x^n:\varphi_1^{(n)}(x^n)=s \\ p_{X^n}(x^n)\leq(1/M_1)\mathrm{e}^{-n\eta} \\ \times p_{X^n|S}(x^n|s)}} p_{X^n}(x^n) \\
&\leq \frac{1}{M_1}\mathrm{e}^{-n\eta} \sum_{s\in\mathcal{M}_1} \sum_{\substack{x^n:\varphi_1^{(n)}(x^n)=s \\ p_{X^n}(x^n)\leq(1/M_1)\mathrm{e}^{-n\eta} \\ \times p_{X^n|S}(x^n|s)}} p_{X^n|S}(x^n|s) \\
&\leq \frac{1}{M_1}\mathrm{e}^{-n\eta}|\mathcal{M}_1| = \mathrm{e}^{-n\eta}.
\end{aligned}
$$

Finally we prove the fourth inequality. We first observe that

$$
p_S(s) = \sum_{x^n:\varphi_1^{(n)}(x^n)=s} p_{X^n}(x^n),\ p_{X^n|S}(x^n|s) = \frac{p_{X^n}(x^n)}{p_S(s)}.
$$

We have the following chain of inequalities:

$$
\begin{aligned}
&p_{SX^nY^n}\left(\mathcal{D}_n^{\mathrm{c}}\cap\mathcal{E}_n\right) \\
&= \sum_{s\in\mathcal{M}_1} p_S(s) \sum_{x^n:\varphi_1^{(n)}(x^n)=s} p_{X^n|S}(x^n|s) \\
&\quad\times \sum_{\substack{y^n:\psi^{(n)}(s,\varphi_2^{(n)}(y^n))=y^n \\ p_{Y^n|S}(y^n|s)\leq(1/M_2)\mathrm{e}^{-n\eta}}} p_{Y^n|X^n}(y^n|x^n) \\
&= \sum_{s\in\mathcal{M}_1} p_S(s) \sum_{\substack{y^n:\psi^{(n)}(s,\varphi_2^{(n)}(y^n))=y^n \\ p_{Y^n|S}(y^n|s)\leq(1/M_2)\mathrm{e}^{-n\eta}}} p_{Y^n|S}(y^n|s) \\
&\leq \sum_{s\in\mathcal{M}_1} p_S(s)\frac{1}{M_2}\mathrm{e}^{-n\eta}\left|\left\{y^n : \psi^{(n)}(s,\varphi_2^{(n)}(y^n)) = y^n\right\}\right| \\
&\overset{\text{(a)}}{\leq} \sum_{s\in\mathcal{M}_1} p_S(s)\frac{1}{M_2}\mathrm{e}^{-n\eta}M_2 = \mathrm{e}^{-n\eta}.
\end{aligned}
$$

Step (a) follows from that the number of y^n correctly decoded does not exceed M_2. ■

Proof of Lemma 1: By definition we have

$$
\begin{aligned}
&p_{SX^nY^n}\left(\mathcal{A}_n\cap\mathcal{B}_n\cap\mathcal{C}_n\cap\mathcal{D}_n\right) \\
&= p_{SX^nY^n}\left\{\frac{1}{n}\log\frac{\hat{q}_{SX^nY^n}(S,X^n,Y^n)}{p_{SX^nY^n}(S,X^n,Y^n)} \geq -\eta,\right. \\
&\qquad \geq -\eta, \\
&\qquad 0 \geq \frac{1}{n}\log\frac{q_{X^n}(X^n)}{p_{X^n}(X^n)} - \eta, \\
&\qquad \frac{1}{n}\log M_1 \geq \frac{1}{n}\log\frac{p_{X^n|S}(X^n|S)}{p_{X^n}(X^n)} - \eta, \\
&\qquad \left.\frac{1}{n}\log M_2 \geq \frac{1}{n}\log\frac{1}{p_{Y^n|S}(Y^n|S)} - \eta\right\}.
\end{aligned}
$$

Then for any $(\varphi_1^{(n)},\ \varphi_2^{(n)},\psi^{(n)})$ satisfying

$$
\frac{1}{n}\log||\varphi_i^{(n)}|| \leq R_i, i = 1,2,
$$

we have

$$
\begin{aligned}
&p_{SX^nY^n}\left(\mathcal{A}_n\cap\mathcal{B}_n\cap\mathcal{C}_n\cap\mathcal{D}_n\right) \\
&\leq p_{SX^nY^n}\left\{\frac{1}{n}\log\frac{\hat{q}_{SX^nY^n}(S,X^n,Y^n)}{p_{SX^nY^n}(S,X^n,Y^n)} \geq -\eta,\right. \\
&\qquad 0 \geq \frac{1}{n}\log\frac{q_{X^n}(X^n)}{p_{X^n}(X^n)} - \eta, \\
&\qquad R_1 \geq \frac{1}{n}\log\frac{p_{X^n|S}(X^n|S)}{p_{X^n}(X^n)} - \eta, \\
&\qquad \left.R_2 \geq \frac{1}{n}\log\frac{1}{p_{Y^n|S}(Y^n|S)} - \eta\right\}.
\end{aligned}
$$

Hence, it suffices to show

$$
\begin{aligned}
\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)}) &\leq p_{SX^nY^n}\left(\mathcal{A}_n\cap\mathcal{B}_n\cap\mathcal{C}_n\cap\mathcal{D}_n\right) \\
&\quad +4\mathrm{e}^{-n\eta}
\end{aligned}
$$

to prove Lemma 1. By definition we have

$$
\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)}) = p_{SX^nY^n}\left(\mathcal{E}_n\right).
$$

Then we have the following.

$$
\begin{aligned}
&\mathrm{P}_{\mathrm{c}}^{(n)}(\varphi_1^{(n)},\varphi_2^{(n)},\psi^{(n)}) = p_{SX^nY^n}\left(\mathcal{E}_n\right) \\
&= p_{SX^nY^n}\left(\mathcal{A}_n\cap\mathcal{B}_n\cap\mathcal{C}_n\cap\mathcal{D}_n\cap\mathcal{E}_n\right) \\
&\quad + p_{SX^nY^n}\left([\mathcal{A}_n\cap\mathcal{B}_n\cap\mathcal{C}_n\cap\mathcal{D}_n]^{\mathrm{c}}\cap\mathcal{E}_n\right) \\
&\leq p_{SX^nY^n}\left(\mathcal{A}_n\cap\mathcal{B}_n\cap\mathcal{C}_n\cap\mathcal{D}_n\right) \\
&\quad + p_{SX^nY^n}\left(\mathcal{A}_n^{\mathrm{c}}\right) + p_{SX^nY^n}\left(\mathcal{B}_n^{\mathrm{c}}\right) + p_{SX^nY^n}\left(\mathcal{C}_n^{\mathrm{c}}\right) \\
&\quad + p_{SX^nY^n}\left(\mathcal{D}_n^{\mathrm{c}}\cap\mathcal{E}_n\right) \\
&\overset{\text{(a)}}{\leq} p_{SX^nY^n}\left(\mathcal{A}_n\cap\mathcal{B}_n\cap\mathcal{C}_n\cap\mathcal{D}_n\right) + 4\mathrm{e}^{-n\eta}.
\end{aligned}
$$

Step (a) follows from Lemma 12. ■

F. *Proof of Lemma 3*

In this appendix we prove Lemma 3.

Proof of Lemma 3: We first prove the following Markov chain (13) in Lemma 3:

$$
SX^{t-1} \leftrightarrow X_t \leftrightarrow Y_t.
$$

We have the following chain of inequalities:

$$
\begin{aligned}
&I(Y_t;SX^{t-1}|X_t)=H(Y_t|X_t)-H(Y_t|SX^{t-1}X_t)\\
&\le H(Y_t|X_t)-H(Y_t|SX^n)\overset{\text{(a)}}{=}H(Y_t|X_t)-H(Y_t|X^n)\\
&\overset{\text{(b)}}{=}H(Y_t|X_t)-H(Y_t|X_t)=0.
\end{aligned}
$$

Step (a) follows from that $S=\varphi_1^{(n)}(X^n)$ is a function of X^n. Step (b) follows from the memoryless property of the information source $\{(X_t,Y_t)\}_{t=1}^{\infty}$. Next we prove the following Markov chain (14) in Lemma 3:

$$
Y^{t-1}\leftrightarrow SX^{t-1}\leftrightarrow (X_t,Y_t).
$$

We have the following chain of inequalities:

$$
\begin{aligned}
&I(X_tY_t;Y^{t-1}|SX^{t-1})\\
&=H(Y^{t-1}|SX^{t-1})-H(Y^{t-1}|SX^{t-1}X_tY_t)\\
&\le H(Y^{t-1}|X^{t-1})-H(Y^{t-1}|X^nSY_t)\\
&\overset{\text{(a)}}{=}H(Y^{t-1}|X^{t-1})-H(Y^{t-1}|X^nY_t)\\
&\overset{\text{(b)}}{=}H(Y^{t-1}|X^{t-1})-H(Y^{t-1}|X^{t-1}Y_t)=0.
\end{aligned}
$$

Step (a) follows from that $S=\varphi_1^{(n)}(X^n)$ is a function of X^n. Step (b) follows from the memoryless property of the information source $\{(X_t,Y_t)\}_{t=1}^{\infty}$. ■

G. *Proof of Lemma 6*

In this appendix we prove Lemma 6.

Proof of Lemma 6 : By the definition of $p^{(\bar{\alpha},\mu,\lambda)}_{SX^tY^t}(s,x^t,y^t)$, for $t=1,2,\cdots,n$, we have

$$
\begin{aligned}
&p^{(\alpha,\mu,\lambda)}_{SX^tY^t}(s,x^t,y^t)\\
&=C_t^{-1}p_{SX^tY^t}(s,x^t,y^t)\\
&\quad\times\prod_{i=1}^{t}f^{(\alpha,\mu,\lambda)}_{q_{X_i}||p_{X_i},p_{X_iY_i|U_i}}(x_i,y_i|u_i).
\end{aligned}
\tag{65}
$$

Then we have the following chain of equalities:

$$
\begin{aligned}
&p^{(\alpha,\mu,\lambda)}_{SX^tY^t}(s,x^t,y^t)\\
&\overset{\text{(a)}}{=}C_t^{-1}p_{SX^tY^t}(s,x^t,y^t)\\
&\qquad\times\prod_{i=1}^{t}f^{(\alpha,\mu,\lambda)}_{q_{X_i}||p_{X_i},p_{X_iY_i|U_i}}(x_i,y_i|u_i)\\
&=C_t^{-1}p_{SX^{t-1}Y^{t-1}}(s,x^{t-1},y^{t-1})\\
&\qquad\times\prod_{i=1}^{t-1}f^{(\alpha,\mu,\lambda)}_{q_{X_i}||p_{X_i},p_{X_iY_i|U_i}}(x_i,y_i|u_i)\\
&\qquad\times p_{X_tY_t|SX^{t-1}Y^{t-1}}(x_t,y_t|s,x^{t-1},y^{t-1})\\
&\qquad\times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t,y_t|u_t)\\
&\overset{\text{(b)}}{=}C_t^{-1}C_{t-1}p^{(\alpha,\mu,\lambda)}_{SX^{t-1}Y^{t-1}}(s,x^{t-1},y^{t-1})\\
&\qquad\times p_{X_tY_t|SX^{t-1}Y^{t-1}}(x_t,y_t|s,x^{t-1},y^{t-1})\\
&\qquad\times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t,y_t|u_t)\\
&=(\Phi_t^{(\alpha,\mu,\lambda)})^{-1}p^{(\alpha,\mu,\lambda)}_{SX^{t-1}Y^{t-1}}(s,x^{t-1},y^{t-1})\\
&\qquad\times p_{X_tY_t|SX^{t-1}Y^{t-1}}(x_t,y_t|s,x^{t-1},y^{t-1})\\
&\qquad\times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t,y_t|u_t).
\end{aligned}
\tag{66}
$$

Steps (a) and (b) follow from (65). From (66), we have

$$
\Phi_t^{(\alpha,\mu,\lambda)}p^{(\alpha,\mu,\lambda)}_{SX^tY^t}(s,x^t,y^t) \tag{67}
$$

$$
\begin{aligned}
&=p^{(\alpha,\mu,\lambda)}_{SX^{t-1}Y^{t-1}}(s,x^{t-1},y^{t-1})\\
&\qquad\times p_{X_tY_t|SX^{t-1}Y^{t-1}}(x_t,y_t|s,x^{t-1},y^{t-1})\\
&\qquad\times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t,y_t|u_t).
\end{aligned}
\tag{68}
$$

Taking summations of (67) and (68) with respect to s,x^t,y^t, we obtain

$$
\begin{aligned}
&\Phi_t^{(\alpha,\mu,\lambda)}\\
&=\sum_{s,x^t,y^t}p^{(\alpha,\mu,\lambda)}_{SX^{t-1}Y^{t-1}}(s,x^{t-1},y^{t-1})\\
&\qquad\times p_{X_tY_t|SX^{t-1}Y^{t-1}}(x_t,y_t|s,x^{t-1},y^{t-1})\\
&\qquad\times f^{(\alpha,\mu,\lambda)}_{q_{X_t}||p_{X_t},p_{X_tY_t|U_t}}(x_t,y_t|u_t),
\end{aligned}
$$

completing the proof. ■

Acknowledgements

We are very grateful to Dr. Shun Watanabe and Dr. Shigeaki Kuzuoka for their helpful comments.

REFERENCES

[1] R. F. Ahlswede and J. Körner, "Source coding with side information and a converse for degraded broadcast channels," *IEEE Trans. Inform. Theory*, vol. IT-21, pp. 629-637, Nov. 1975.

[2] A. D. Wyner, "On source coding with side information at the decoder," *IEEE Trans. Inform. Theory*, vol. IT-21, pp. 294-300, May 1975.

[3] R. Ahlswede, P. Gàs, and J. Körner, "Bounds on conditional probabilities with applications in multi-user communication," *Z. Wahrscheinlichkeitstheorie verw. Gebiete*, vol. 34, pp. 157-177, 1976.

[4] A. D. Wyner and J. Ziv, "The rate-distortion function for source coding with side information at the decoder," *IEEE Trans. Inform. Theory*, vol. IT-22, pp. 1-10, Jan. 1976.

[5] W. Gu and M. Effors, "A strong converse for a collection of network source coding problems," *Proceedings of the IEEE International Symposium on Information Theory*, Soul, Korea, pp.2356-2320, June 2009.

[6] D. Slepian and J. K. Wolf, "Noiseless coding of correlated information sources," *IEEE Trans. Inform. Theory*, vol. IT-19, pp. 471-480, July 1973.

[7] Y. Oohama and T. S. Han, "Universal coding for the Slepian-wolf data compression system and the strong converse theorem,"*IEEE Transactions on Information Theory*, vol IT-40, no.6, 1908-1919, Nov. 1994.

[8] I. Csiszár and J. Körner, *Information Theory: Coding Theorems for Discrete Memoryless Systems.* London, U.K., Academic, 1981.

[9] T. S. Han, *Information-Spectrum Methods in Information Theory.* Springer-Verlag, Berlin, New York, 2002. The Japanese edition was published by Baifukan-publisher, Tokyo, 1998.